

***Six2* EXHIBITS A TEMPORAL-SPATIAL EXPRESSION PROFILE
IN THE DEVELOPING MOUSE PALATE AND IMPACTS CELL
PROLIFERATION DURING MURINE PALATOGENESIS**

**A Thesis Submitted to the College of Graduate Studies and
Research in Partial Fulfilment of the Requirements for the Degree
of Master of Science in the College of Pharmacy and Nutrition
University of Saskatchewan
Saskatoon**

By Dennis Okori Okello

© Copyright Dennis Okori Okello, July 2015. All rights reserved.

PERMISSION TO USE

In presenting this thesis in partial fulfilment of the requirements for a Master of Science degree from the University of Saskatchewan, I agree that the Libraries of this University may make it freely available for inspection. I also agree that permission for copying this thesis in any manner, in whole or in part, for scholarly purposes may be granted by Dr. Adil J. Nazarali, the Professor who supervised my thesis work, or in his absence, by the Dean of the College of Pharmacy and Nutrition in which my thesis work was done. It is understood that any copying or publication, or use of this thesis or parts thereof for financial gain shall not be allowed without my written permission. It is also understood that due recognition shall be given to me and to the University of Saskatchewan in any scholarly use which may be made of any material in my thesis.

Requests for permission to copy or to make use of material in this thesis in whole or in part should be addressed to:

Dean of the College of Pharmacy and Nutrition
University of Saskatchewan
104 Clinic Place
Saskatoon, Saskatchewan
S7N 2Z4, CANADA.

ABSTRACT

Cleft palate is one of the most common congenital malformations in humans which occurs at a frequency of approximately 1:700 live births worldwide. *Sine Oculis-related homeobox 2* (*Six2*) is a member of the vertebrate *Six* gene family that encode proteins that are transcription factors. *Six2* has been reported to be a downstream target of *Homeobox a2* (*Hoxa2*), a gene that plays a direct a role in mouse secondary palate (SP) development.

In my thesis, I utilized quantitative real time Polymerase Chain Reaction (qPCR), Western blot analysis and fluorescence immunohistochemisrty (IHC) to characterize the spatial and temporal distribution patterns of *Six2* in the developing SP. Additionally, I also employed *in vivo* cell counting analysis and *in vitro* cell proliferation assays to investigate the role of *Six2* during palate mesenchymal cell proliferation.

My study examined the temporal and spatial distribution of *Six2* in the developing mouse palatal mesenchyme and epithelia in both wild-type and *Hoxa2* null mice. *Six2* was expressed throughout the period of embryonic palatogenesis, with the highest levels of *Six2* mRNA and protein observed in palatal shelves at E13.5 in both wild-type and *Hoxa2* null mice. *Six2* protein expression at all stages of SP development (E12.5 to E15.5) increased in the anterior to posterior (A-P) direction with highest expression in the posterior regions of the developing SP. In addition, expression of *Six2* protein was higher in the oral half of the palatal mesenchyme compared to the nasal half of the palatal mesenchyme. Interestingly, *Six2* protein was expressed in the nasal palatal epithelium but was completely absent from the oral palatal epithelium. Loss of the *Hoxa2* gene induced up regulation of *Six2* protein and mRNA in the developing palate across all stages of palatogenesis. In the *Hoxa2* null mice, there was a significant increase in cell proliferation (Ki-67 positive cells) and

the percentage of actively proliferating cells that were co-expressing *Six2* protein (*Six2*/Ki-67 double positive cells) along both the A-P and oral-nasal (O-N) axes of the developing SP. Also, the highest percentage of actively proliferating cells and *Six2*/Ki-67 double positive cells was observed in the nasal half of the posterior palatal mesenchyme. Furthermore, *Six2* siRNA knock down in mouse embryonic palatal mesenchyme (MEPM) cell cultures restored cell proliferation and *Cyclin D1* expression in the *Hoxa2* null cell cultures to wild-type levels.

Collectively, my data reveals a novel spatial and temporal expression profile for *Six2* in the developing mouse SP and the potential role it might play during the epithelial-mesenchymal cross talk that drives palatal shelf cell proliferation and out growth.

ACKNOWLEDGEMENTS

I would like to sincerely thank my supervisor Dr. Adil J. Nazarali for the never ending support and encouragement that he offered during my programme. This work would not have been possible without his support. My sincere gratitude also goes to Dr. William Kulyk for the insightful advice he provided during my graduate committee meetings and during the editing of this thesis. I would also like to thank in a special way Dr. Ed Krol, the Graduate Chair of Pharmacy, in the College of Pharmacy and Nutrition for always finding time to chair my committee meetings. Collectively, these three individuals provided the advice and suggestions that helped guide my research project. I would also like to thank my laboratory colleagues; Paul, Muath, Ran, Merlin, and Jotham with whom I shared the highs and lows of graduate student life.

Finally, I would like to specially thank the College of Pharmacy and Nutrition of the University of Saskatchewan for funding part of my studies.

DEDICATION

This Degree is dedicated to my supportive family especially my Dad, Dr. John Okorio, without whom I would never have dreamt of coming to Canada to pursue graduate studies. I will forever be indebted to you for your support both financially and morally, that you have rendered me in my never-ending pursuit for knowledge. Thank you for believing in me.

TABLE OF CONTENTS

PERMISSION TO USE	i
ABSTRACT	ii
ACKNOWLEDGEMENTS	iv
DEDICATION.....	v
TABLE OF CONTENTS	vi
LIST OF FIGURES.....	ix
LIST OF ABBREVIATIONS	xi
1. INTRODUCTION.....	1
2. REVIEW OF LITERATURE	3
2.1 An overview of craniofacial development	3
2.1.1 Early embryonic development	3
2.1.2 Migration of the cranial neural crest cell population	6
2.1.3 Formation of the face	12
2.2 Secondary palate development	15
2.3 Molecular signalling in the developing palate	17
2.4 Molecular signalling along the A-P axis of the developing SP	20
2.5 Molecular signalling along the O-N axis of the developing SP	22
2.6 The <i>Six</i> family of genes.....	23
2.7 Expression patterns of <i>Six</i> genes	24
2.8 <i>SIX</i> genes and genetic diseases	28
2.9 The <i>Six2</i> gene	30
3. HYPOTHESES	32
4. OBJECTIVES	32
5. MATERIALS AND METHODS	33
5.1 <i>Hoxa2</i> transgenic mice	33
5.2 PCR analysis of genotype	33
5.3 Immunohistochemistry (IHC)	34

5.4 RNA Isolation and Reverse Transcription	35
5.5 Quantitative Real Time RT-PCR	36
5.6 Droplet digital PCR (ddPCR)	36
5.7 Western blot analysis.....	37
5.8 Culturing MEPM cells.....	39
5.9 siRNA treatment and cell proliferation analysis	40
5.10 Statistical analysis	41
6. RESULTS	42
6.1 Six2 protein exhibits a temporal distribution pattern in the developing secondary palate	42
6.2 Six2 protein is spatially distributed in both the epithelium and mesenchyme of the developing secondary palate	44
6.3 Six2 protein exhibits an A-P differential expression pattern in the developing secondary palate	47
6.4 Six2 protein and mRNA are up-regulated in the absence of <i>Hoxa2</i>	50
6.5 Analysis of cell proliferation in vivo.....	54
6.6 Analysis of cell proliferation in vitro.....	57
7. DISCUSSION	63
7.1 <i>Six2</i> exhibits a temporal-spatial expression profile in the developing secondary palate	63
7.2 Six2 protein exhibits an A-P differential expression pattern in the developing secondary palate.	67
7.3 <i>Six2</i> mRNA and protein are up-regulated in the absence of <i>Hoxa2</i>	68
7.4 <i>Six2</i> has an impact on palate mesenchymal cell proliferation during palatogenesis.....	71
7.4.1 There is increased cell proliferation and a higher percentage of cells co-expressing Six2 and Ki-67 in the nasal half of the posterior palatal mesenchyme of <i>Hoxa2</i> null palatal shelves.	71
7.4.2 In vitro <i>Six2</i> siRNA knockdown restored cell proliferation and <i>Cyclin D1</i> expression in <i>Hoxa2</i> null MEPM cell cultures to wild-type levels.....	73

8. CONCLUSION	75
9. FUTURE WORK CONSIDERATIONS.....	77
10. REFERENCES.....	81
11. APPENDIX.....	96
In vivo cell counting analysis.....	96
Absolute quantitative expression of palatal <i>Six2</i> mRNA by droplet digital PCR.....	104
Sample of genotyping results of <i>Hoxa2</i> transgenic mice.....	105
Primers used for qPCR analysis of <i>Msx1</i> , <i>Bmp4</i> , <i>Barx1</i> and <i>Ptx1</i>	106
An agarose gel showing the integrity of the primers that were used for PCR genotyping analysis	107
Reagent catalogue numbers	108
Letters of permission to use images.....	109

LIST OF FIGURES

Figure 1: A schematic illustration of an early embryo showing the prosencephalon, mesencephalon and rhombencephalon.....	5
Figure 2: A schematic representation of early embryonic development.....	8
Figure 3: A diagrammatic illustration of cranial neural crest cell migration.....	10
Figure 4: A schematic demonstration of border induction and neurulation.....	11
Figure 5: A diagrammatic illustration of facial development.....	14
Figure 6: A schematic illustration of secondary palate development.....	16
Figure 7: A schematic illustration of the human <i>SIX2</i> gene structure.....	26
Figure 8: A schematic drawing of embryonic expression patterns and chromosomal locations of mouse <i>Six</i> family genes.....	27
Figure 9: <i>Six2</i> protein exhibits a temporal distribution pattern in the developing palate.....	43
Figure 10: <i>Six2</i> protein exhibits a spatial expression pattern in the palate epithelium.....	45
Figure 11: <i>Six2</i> protein exhibits a spatial expression pattern in the palatal mesenchyme.....	46
Figure 12: <i>Six2</i> protein exhibits an Anterior-Posterior differential expression pattern in the palate epithelium and mesenchyme.....	48
Figure 13: <i>Six2</i> mRNA and protein are up-regulated in the absence <i>Hoxa2</i>	51
Figure 14: <i>In vivo</i> cell counting analysis of <i>Six2</i> protein positive cells in the nasal half of the palatal mesenchyme in wild-type and <i>Hoxa2</i> null littermates showing an expanded domain of <i>Six2</i> protein expression in the <i>Hoxa2</i> null palatal shelves.....	53

Figure 15: Increased cell proliferation in nasal half of the posterior palatal mesenchyme of <i>Hoxa2</i> null palatal shelves.....	55
Figure 16: The number of cells co-expressing <i>Six2</i> and <i>Ki-67</i> is highest in the nasal half of the posterior palatal mesenchyme of <i>Hoxa2</i> null palatal shelves.....	56
Figure 17: <i>Six2</i> siRNA knockdown in MEPM cells.....	59
Figure 18: <i>In vitro</i> <i>Six2</i> siRNA knock down restored cell proliferation and <i>Cyclin D1</i> expression in the <i>Hoxa2</i> null palate mesenchymal cell cultures to wild-type levels.....	60
Figure 19: <i>In vitro</i> <i>Six2</i> siRNA knock-down did not have an effect on the expression of <i>Msx1</i> , <i>Bmp4</i> , <i>Barx1</i> and <i>Ptx1</i>	62
Figure 20: A schematic diagram illustrating the spatial expression pattern of <i>Six2</i> protein in the developing palate at E13.5 and E14.5.....	66
Figure 21: A schematic diagram demonstrating an expanded domain of <i>Six2</i> protein expression in the mesenchyme of the <i>Hoxa2</i> null palatal shelf at E14.5.....	70
Figure 22: A schematic illustration of the pathways involved in Oral-Nasal patterning during palatal shelf out growth.....	79

LIST OF ABBREVIATIONS

ANOVA	Analysis of variance
A-P	Anterior-Posterior
BA	Branchial Arch
Barx1	BarH-like homeobox 1
Bmp4	Bone morphogenetic protein 4
bp	base pair
C	Celsius
cDNA	complementary DNA
ChIP	Chromatin immunoprecipitation
CL/P	Cleft lip and palate
CNCC	Cranial neural crest cell
CO ₂	Carbon dioxide
CP	Cleft Palate
Ct	Threshold cycle
ddPCR	Droplet digital Polymerase Chain Reaction
Dlx5	Distal-less homeobox 5
DM	Myotonic dystrophy
DMEM	Dulbecco's modified eagle's medium
DMPK	Myotonic dystrophy protein kinase
DMWD	Myotonic dystrophy with beta-transducin (WD) repeat
DNA	Deoxyribonucleic acid
E	Embryonic day
EDTA	Ethylenediaminetetraacetic acid
Eya1	Eyes absent1
Fgf	Fibroblast growth factor
Fgf10	Fibroblast growth factor 10
Fgfr	Fibroblast growth factor receptor
FIG	Figure
Foxf1a	Forkhead box F1a
Foxf2	Forkhead box F2
h	Hour
HD	Homeodomain
Hoxa2	Homeobox a2
HPE	Holoprosencephaly
IDV	Integrated density value
IgG	Immunoglobulin G
IHC	immunohistochemistry/immunohistochemical
<i>Italics</i>	Denotes the gene or mRNA form

M	Molar
Meox2	Mesenchyme homeobox 2
MEPM	Mouse embryonic palatal mesenchyme
MES	Midline Epithelial Seam
mg	milligrams
min	Minutes
mM	Millimolar
Mn1	Meningioma 1
MNP	Medial nasal prominence
mRNA	Messenger Ribonucleic acid
Msx1	msh-like1
MXP	Maxillary prominence
n	Number of samples
NaCl	Sodium chloride
NCC	Neural crest cell
ng	Nanogram
nM	Nanomolar
°C	Degrees centigrade
OCT	Optimal cutting temperature
O-N	Oral-Nasal
Osr1	Odd-skipped related 1
Osr2	Odd-skipped related 2
Pax9	Paired box gene 9
PBS	Phosphate buffered saline
PCR	Polymerase Chain Reaction
PP	Primary Palate
Ptc	Patched 1
Ptx1	Paired like homeodomain transcription factor1
PVDF	Polyvinylidene difluoride
RIPA	Radio immuno precipitation assay buffer
RQ	Relative quantification
RT	Reverse transcription
RTM	Room temperature
RT-PCR	Reverse Transcriptase Polymerase Chain Reaction
s	Seconds
SD	Six domain
Shh	Sonic hedgehog
Shox2	Short stature homeobox 2
siRNA	Small interfering ribonucleic acid
Six	Sine Oculis-related homeobox

Six2	Sine Oculis-related homeobox 2
Smo	Smoothened
SM-PBS	Skim milk in phosphate buffered saline
SM-PBST	Skim milk in phosphate buffered saline with 1% Triton X-100
SP	Secondary palate
Spry2	Sprouty homolog 2
Tbx22	T-box transcription factor 22
Tgf β	Transforming growth factor beta
TGif	TG-interacting factor
UTR	Untranslated region
UV	Ultraviolet
Wt	Wild-type
μ l	Microlitre
μ m	Micrometer
μ M	Micromolar

1. INTRODUCTION

Cleft palate is one of the most common congenital malformations in humans, occurring at a frequency of approximately 1:700 live births worldwide (Gorlin et al., 2001). The palate separates the nasal and oral cavities and this allows for respiration, feeding and phonation to occur. Environmental and genetic factors have both been implicated in the pathogenesis of the cleft palate. Maternal use of the anticonvulsant drugs valproic acid and phenytoin, has been shown to cause cleft palate in the newborn (Holmes et al., 2001). However, the molecular mechanisms involved in this drug-induced manifestation are poorly understood.

Craniofacial development is a tightly regulated process that involves the growth and fusion of the maxillary, mandibular and nasal processes (Chai and Maxson, 2006). Mouse secondary palate development begins around E11.5. At E12.0-13.5, paired palatal shelves initiate from the oral side of the maxillary processes and grow vertically downwards on either side of the tongue. The tongue depresses at E14.0 causing the palatal shelves to reorient above the tongue. These then grow horizontally towards each other ultimately touching at E14.5 to form the midline epithelial seam (MES). The MES degrades by E15.5, leaving behind a confluent secondary palate. In addition, the secondary palate fuses anteriorly with the primary palate, a derivative of the medial nasal processes, to form the intact roof of the oral cavity. Therefore, disruption of the growth, elevation or fusion of the palatal shelves can lead to cleft palate (Bush and Jiang, 2012; Ferguson, 1988; Gritli-Linde, 2007, Smith et al., 2013).

Six2 is a member of the vertebrate gene family consisting of the following genes: *Six1*, *Six3*, *Six4*, *Six5* and *Six6* which encode transcription factor proteins homologous to the *Drosophila* 'sine oculis' homeobox protein (Kiyoshi et al., 2000). *Six2* has been shown to be involved in regulating

craniofacial and kidney development (Fogelgren et al., 2008, 2009; Lozanoff, 1993; Lozanoff et al., 2001; Ma and Lozanoff, 1996, 1999; McBratney et al., 2003; Singh et al., 1998). *Six2* expression has been reported in the mesenchyme of the nasal prominences of E11.5 mice (Brodbeck et al., 2004; Fogelgren et al., 2008; Ohto et al., 1998; Oliver et al., 1995). *Six2* has also been reported to be a direct downstream target of *Hoxa2* in the murine branchial arches (Kutejova et al., 2005; Kutejova et al., 2008).

Although *Six2* has been reported to be expressed during mammalian craniofacial development, no detailed expression analysis has been performed in the developing secondary palate. In addition, its precise role in the craniofacial region and especially in the developing mammalian secondary palate is still unknown. My thesis therefore seeks to address these gaps in knowledge regarding the transcription factor *Six2* during palatogenesis.

2. REVIEW OF LITERATURE

2.1 An overview of craniofacial development

2.1.1 Early embryonic development

In the human embryo, oral development starts early by the appearance of the prechordal plate in the bilaminar germ disc on the 14th day of development. The endodermal thickening of the prechordal plate demarcates the cranial pole of the oval embryonic disk, which later contributes to the oropharyngeal membrane. This bilaminar membrane is the site of the junction of the ectoderm that forms the mucosa of the mouth and the endoderm that forms the mucosa of the pharynx, which is the most cranial part of the foregut. The oropharyngeal membrane is one of two sites of contact between ectoderm and endoderm, where mesoderm fails to intervene between the two primary germ layers; the other site is the cloacal membrane at the terminal end of the hindgut. The oropharyngeal membrane also demarcates the stomodeum, the primitive mouth that forms the center of the developing face (Richman et al., 1995; Sulik et al., 1990).

The development of the head relies on the inductive activities of the prosencephalic and rhombencephalic organizing centres that are regulated by *sonic hedgehog* (*Shh*) (Hu et al., 1999) (Fig. 1). Located at the rostral end of the notochord beneath the forebrain (prosencephalon) is the prosencephalic centre which is derived from the prechordal mesoderm that migrates from the primitive streak. This centre induces the visual and inner-ear apparatus and the upper third of the face through *Shh* protein signalling. Similar to the prosencephalic centre, the caudal rhombencephalic centre located beneath the hindbrain (Rhombencephalon) also induces the middle and lower thirds of the face including the middle and external ears through *Shh* signalling. In

association with these developments is the bifurcation of the unilobar forebrain (prosencephalon) to form paired telencephalic hemispheres and evagination of the paired olfactory bulbs and optic nerves. Failure of these cerebral divisions leads to severe facial anomalies (Johnson et al., 1995).




Figure 1: A schematic illustration of showing the three primary brain vesicles of an early embryo (prosencephalon, mesencephalon and rhombencephalon) and their five main subdivisions.

(<http://what-when-how.com/neuroscience/development-of-the-nervous-system-gross-anatomy-of-the-brain-part-1/>. Web. October 4th 2014).

2.1.2 Migration of the cranial neural crest cell population

Formation of cranial neural crest cells (CNCC) is one of the key features of craniofacial development. The specification, emigration and migration, proliferation, survival and ultimate fate determination of the CNCC play an important role in regulating craniofacial development. CNCC give rise to a variety of cell types during embryonic development. These cells eventually give rise to the hard tissues of the head such as bone, cartilage and teeth; optic elements such as the choroid, sclera, iris, and ciliary body, and sensory ganglia of the fifth, seventh, ninth and tenth cranial nerves (Noden, 1983; Couly et al., 1993; Kontges and Lumsden, 1996) (Fig. 2). Genetic disorders, environmental insults, or a combination of both can alter the fate determination of CNCC leading to craniofacial malformations.

CNCC arise uniformly at the dorso-lateral edge of the closing neural folds, along almost the entire length of the vertebrate embryo neural axis. This region corresponds to the interface between the non-neural ectoderm (presumptive epidermis or surface ectoderm) and the neural plate (neuroepithelium), a region that is commonly referred to as the neural plate border. CNCC induction requires contact mediated interactions between the surface ectoderm and neuroepithelium and each of these tissues contributes to the neural crest cell lineage (Selleck and Bronner-Fraser, 1995). In addition to their induction along the dorsolateral edge of the neural plate, neural crest cells undergo an epithelial to mesenchymal transformation whereby they delaminate from the neural tube and begin to migrate (Figs. 2, 3). These epithelial-mesenchymal transitions are marked by changes in cell adhesion and cytoarchitecture (Bronner-Fraser, 1993; LaBonne and Bronner-Fraser, 1999; Le Douarin et al., 2004). Following delamination from the neural tube, CNCC migrate along specific pathways to their final destinations (Fig. 4). In

vertebrates, the majority of CNCC are derived from the hindbrain, which migrate ventrolaterally from neural tube in three distinct sub-ectodermal streams adjacent to the even numbered rhombomeres (r2, r4, r6). The proliferative activity of the CNCC in the craniofacial region eventually leads to the formation of the frontonasal processes and discrete swellings that demarcate each branchial arch. The three streams of CNCC populate the first, second and third branchial arches respectively in keeping with their craniocaudal axial origins (Lumsden et al., 1991) and give rise to a wide variety of cell lineages, which are distinct for each branchial arch (Noden, 1983; Kontges and Lumsden, 1996).




Figure 2: A schematic representation of early embryonic development. Panels A-D depict early development (at the third and fourth weeks of gestation) in which the neural plate (A), neural groove (B), and neural tube (C) are formed from the dorsal surface of the embryo. The left side of each panel depicts the developing embryo in a dorsal view, and the right side shows cross sections through the nervous system cut at the levels indicated by the arrows. Note also the cells formed from differentiated cells of the neural crest (D).

(Taken from <http://what-when-how.com/neuroscience/development-of-the-nervous-system-gross-anatomy-of-the-brain-part-1/>. Web. October 10th 2014).




Figure 3: Diagrammatic illustration of cranial neural crest cell formation and delamination.

(REECE, JANE B.; URRY, LISA A.; CAIN, MICHAEL L.; WASSERMAN, STEVEN A.; MINORSKY, PETER V.; JACKSON, ROBERT B., CAMPBELL BIOLOGY, 9th Edition, © 2011. Reprinted and Electronically reproduced by permission of Pearson Education, Inc., Upper Saddle River, NJ).




Figure 4: A schematic demonstration of border induction and neurulation. The neural plate border (green) is induced by signalling between the neuroectoderm (purple) and the non-neural ectoderm (blue) and from the underlying paraxial mesoderm (yellow). During neurulation, the neural plate borders (neural folds) elevate, causing the neural plate to roll into a neural tube. Neural crest cells (green) delaminate from the neural folds or the dorsal neural tube (shown), depending on the species and axial level. (Reprinted by permission from Macmillan Publishers Ltd: Nature Reviews Neuroscience., Neuro crest specification: migrating into genomics., copyright Oct 1, 2003.)

2.1.3 Formation of the face

The face originates from five prominences that surround a central depression, the stomodeum, which constitutes the future mouth. These prominences include the single median frontonasal and the paired maxillary and mandibular prominences ; the latter two are derivatives of the first pair of six branchial arches. These prominences and arches arise in most part, from neural crest ectomesenchyme (neural crest cells) that migrates from the dorsal neural tube (point of fusion of the neural plate borders) into the facial and neck regions (Fig. 2).

The frontonasal prominence surrounds the forebrain, which has lateral optic diverticula that form the eyes. The frontal portion of the prominence between the eyes forms the forehead; at the inferolateral corners, thickened ectodermal nasal (olfactory) placodes arise. These placodes become the olfactory epithelium and develop the underlying olfactory nerves. Union of the facial prominences occurs by merging of the frontonasal, maxillary, and mandibular prominences. Merging of what are initially incompletely separated prominences occurs as the intervening grooves disappear as a result of migration into and proliferation of underlying mesenchyme in the groove. Fusion of the freely projecting medial nasal prominences with the maxillary and lateral nasal prominences on each side requires the disintegration of their contacting surface epithelia, allowing the underlying mesenchymal cells to intermingle. Failure of normal disintegration by cell death or mesenchymal transformation is a cause of cleft upper lip and primary palate, as such failure prevents the intermingling of maxillary and medial nasal mesenchyme. Fusion of the medial nasal and maxillary prominences provides for continuity of the upper jaw and lip and separation of the nasal pits from the stomodeum. The midline merging of the medial nasal prominences forms the median

tuberculum and philtrum of the upper lip, the tip of the nose, and the primary palate.

The intermaxillary segment of the upper jaw (the premaxilla), in which the four upper incisor teeth will develop, arises from the median primary palate that is initially a widely separated pair of inwardly projecting swellings of the merged medial nasal prominences. Abnormal unilateral clefting resulting from failed fusion of medial nasal and maxillary prominences produces a deflection of the nose and upper lip.

The lower jaw and lip are very simply formed by midline merging of the paired mandibular prominences and are the first parts of the face to become definitively established. The lateral merging of the maxillary and mandibular prominences creates the commissures (corners) of the mouth (Fig. 5) (Gilbert, 2003; Lumsden and Keynes, 1989; Helms et al., 2005; Sperber and Sperber, 2009; Hinrichsen, 1985).




Figure 5: A diagrammatic illustration of facial development Showing the prominences from which the structures of the face are formed. The frontonasal or median nasal prominence (red) gives rise to the forehead (A), centre of the nose (B), philtrum of the upper lip (C) and the primary palate (D). The lateral nasal prominences (blue) form the sides of the nose (B,D). The maxillary and mandibular prominences (green) give rise to the sides of the middle and lower face, lateral borders of the lips, the secondary palate (specifically from maxillary prominences) and to the lower jaw (specifically from the mandibular prominences) (D).

(REPRODUCED WITH PERMISSION FROM HELMS JA, CORDERO D, TAPADIA MD. (2005). NEW INSIGHTS INTO CRANIOFACIAL MORPHOGENESIS. DEVELOPMENT 132:851–861).

2.2 Secondary palate development

The primary and secondary palates form the main barrier between mammalian nasal and oral cavities. This was an evolutionary advancement from the single common chamber seen in birds and reptiles (Chong et al., 2002 and Sperber et al., 2002). The secondary palate (SP) is the primordium of both the hard and soft palates of the palate. SP development is a multi-step process involving palatal shelf growth, elevation, midline fusion of the two palatal shelves and the disappearance of the midline epithelial seam (MES). The palatal structures are composed of the cranial neural crest (CNC-derived ectomesenchyme and pharyngeal ectoderm) (Ferguson, 1988; Shuler, 1995).

Palatal shelves are covered by epithelia that can be divided into the nasal, oral and medial edge epithelium (MEE). The nasal and oral epithelia differentiate into pseudostratified columnar and stratified squamous epithelia respectively, whereas the medial edge epithelium gets removed by means of programmed cell death (apoptosis) and cell migration (Martinez-Alvarez et al., 2000; Vaziri Sani et al., 2005).

Secondary palate development begins at E11.5 in mice and around the sixth week of gestation in humans. Initially, the palatal shelves exist as projections in the internal aspects of the maxillae, on either side of the tongue. These two shelves then grow vertically downwards on either sides of the tongue between the stages E12.0 and E13.5 (Ferguson, 1988; Berkovitz, et al., 2002). As the jaws develop, the tongue depresses causing the palatal shelves to re-orient themselves above the tongue at E14.0. The shelves then start growing horizontally towards each other where they ultimately touch at E14.5 to form the midline epithelial seam (MES). The MES degrades by E15.5 leaving behind a confluent SP (Fig. 6).




Figure 6: A schematic illustration of secondary palate development showing the palatal shelves emerging from the maxillary prominences at E11.5, growing vertically downwards on either side of the tongue (E12.5), re-orienting above the tongue (E13.5-E14.0), growing horizontally and touching to form the midline epithelial seam (E14.5) and eventually fusing to form a solid, confluent secondary palate (E15.5). ps- Palatal shelf, T-tongue.

(REPRODUCED WITH PERMISSION OF THE AMERICAN SOCIETY OF CLINICAL INVESTIGATION IN THE FORMAT REPUBLISH IN A THESIS/DISSERTATION VIA COPYRIGHT CLEARANCE CENTRE).

2.3 Molecular signalling in the developing palate

Mammalian SP development is an intricate process involving the participation of several transcription factors and growth factors that interact with their receptors as well as with each other. These networks work together to regulate critical cellular processes including cell proliferation, migration and epithelial-mesenchymal transdifferentiation and apoptosis. Similar to the development of other organs in the embryo, SP palate development relies on the sequential and reciprocal epithelial-mesenchymal interactions. Some of the transcription factors and growth factors involved in SP development include; *Shh*, *Msx1*, *Bmp4*, *Barx1*, *Fgf10*, *Fgf* receptors, *Pax9*, *Osr2* and *Tgfb*.

Shh is a member of the hedgehog family of secreted proteins which plays crucial roles in diverse developmental processes including left-right axis establishment, dorso-ventral patterning of the neural tube, endoderm development, A-P patterning of the developing limb, brain development and patterning (McMahon et al., 2003; Roessler and Muenke, 2003). *Shh* is expressed at all stages of SP development and follows a striped pattern of expression corresponding to areas of thickened epithelium that later form the palatal rugae. The A-P outgrowth of palatal shelves has been associated with periodic formation of palatal rugae and is controlled by reciprocal epithelial-mesenchymal crosstalk along the A-P axis (Lan and Jiang, 2009; Pantalacci et al., 2008; Welsh and O'Brien, 2009). Palatal rugae have been reported to express high levels of *Shh* in addition to acting as signalling centers involved in epithelial-mesenchymal interactions required to coordinate palate outgrowth and patterning (Lan and Jiang, 2009; Rice et al., 2006; Welsh and O'Brien, 2009). A study by Han et al., 2009 reported that the epithelial cells expressing *Shh* are not actively proliferating, but the mesenchymal cells underlying these regions are more highly proliferative than mesenchymal cells in other areas of

the palate. *Shh* is a downstream target of the *Msx1* network that regulates cell proliferation in the anterior palate (Zhang et al., 2002).

Loss of the *Spry2* gene has been associated with the disorganization in the expression pattern of *Shh*, which ultimately leads to deformities in the rugae in the palate of these knockout animals (Welsh et al., 2007). Studies of mice involving epithelium-specific inactivation of *Shh* or mesenchyme-specific inactivation of *smoothed* (*Smo*), which encodes a transmembrane protein required for transducing *Shh* signalling, have demonstrated that *Shh* signals from the epithelium to the underlying mesenchyme to promote palatal cell proliferation and outgrowth (Rice et al., 2004; Lan and Jiang, 2009). In addition, the exogenous application of *Shh* has been shown to induce a mitogenic response in palatal explant cultures (Zhang et al., 2002; Rice et al., 2004). This mitogenic effect is mediated, in part, by the cell cycle regulators *Cyclin D1* and *Cyclin D2*, the expression of which was reduced in the palatal mesenchyme of embryos with mesenchyme-specific inactivation of *Smo*.

Lan and Jiang, (2009) reported that *Shh* from palatal epithelial cells stimulates cell proliferation and the expression of *Fgf10* in the developing palatal mesenchyme. *Shh* and *Fgf10* have also been reported to function in a positive-feedback loop to drive the outgrowth of the palatal shelves (Bush and Jiang., 2012; Lan and Jiang., 2009; Rice et al., 2004). The expression of *Foxf1a* and *Foxf2* and *Osr2* genes was also reduced in the palatal mesenchyme, indicating that these transcription factors might be downstream effectors of *Shh* signalling (Lan and Jiang, 2009). *Fgf10* is a crucial mesenchymal signal that is required for palatal outgrowth. Mice homozygous for a null mutation in either *Fgf10* or the gene encoding its receptor, *Fgfr2b*, exhibited cleft palate with impaired palatal shelf outgrowth (Rice et al., 2004). Whereas *Fgf10* mRNA expression was restricted to the mesenchyme, *Fgfr2b* mRNA was most

abundantly detected in the overlying epithelium. *Fgfr2* function is required within the epithelium, as mice harbouring an epithelial-specific deletion of *Fgfr2* also exhibited cleft palate (Hosokawa et al., 2009). Both epithelial and mesenchymal cell proliferation were reduced in the absence of either *Fgf10* or *Fgfr2b*, however, suggesting the presence of a factor that signals from the epithelium back to the underlying mesenchyme and that is dependent on *Fgf10/Fgfr2b* signalling. Indeed, *Shh* expression was dramatically reduced in the epithelium of *Fgf10* null and *Fgfr2b* null embryos, implying that the decreased palatal mesenchymal cell proliferation observed in these mutants might be a consequence of reduced *Shh* expression in the epithelium (Rice et al., 2004). The expression of *Fgf10* was also reduced in the palatal mesenchyme of embryos lacking mesenchymal *Smo*, indicating that *Shh* and *Fgf10* function in a positive-feedback loop that drives the outgrowth of the palatal shelves (Lan and Jiang, 2009).

In addition, cross-regulation of the *Shh* and *Bmp* signalling pathways has also been detected in the developing palate (Baek et al., 2011; Lan and Jiang, 2009; Zhang et al., 2002). The loss of *Smo* within the palatal mesenchyme led to up regulation of *Bmp4* and down regulation of *Bmp2* (Lan and Jiang, 2009). The positive regulation of *Bmp2* by *Shh* signalling is consistent with the finding that exogenous *Shh*-containing beads induce *Bmp2* expression in palatal explant culture (Zhang et al., 2002). Whereas complete inactivation of *Bmp4* resulted in early embryonic lethality, ablation of *Bmp4* function in the maxillary mesenchyme and throughout the oral epithelium caused a cleft lip phenotype, but no secondary palate defect was reported (Liu et al., 2005). Overexpression of the *Bmp* antagonist noggin specifically in the palatal mesenchyme led to retarded palatal growth and cleft palate in mice (Xiong et al., 2009), further supporting the involvement of *Bmp* signalling during palatal growth. Studies

have also shown that *Bmp* signalling during palatogenesis occurs via the type I Bmp receptor *Bmpr1a*. Disruption of *Bmpr1a* in the maxillary mesenchyme and throughout the oral epithelium resulted in cleft lip and palate (Liu et al., 2005).

Pax9 is a member of the transcription factor family characterized by the paired-class DNA binding domain (Stapleton et al., 1993). *Pax9*-deficient mice have been reported to die shortly after birth, exhibiting complete cleft palate (Peters et al., 1998). Yu et al., 2013 reported that *Pax9* regulates epithelial-mesenchymal signalling and A-P patterning during palate outgrowth and that *Pax9* acts upstream of *Osr2* to regulate palatal shelf growth and elevation. In addition, this study also reported that *Pax9* acts upstream of *Bmp4*, *Fgf10*, *Msx1* and *Osr2* in the palatal mesenchyme.

2.4 Molecular signalling along the A-P axis of the developing SP

Although the palatal shelves appear as continuous outgrowths that project from the maxillary processes into the oral cavity, studies have revealed differential gene expression along the A-P axis, and that this heterogeneity is evident from the early stages of palatal outgrowth (Hilliard et al., 2005; Li and Ding, 2007; Welsh and O'Brien, 2009; reviewed in Smith et al., 2013). The SP may be divided into three regions along the A-P axis, namely the anterior palate, medial palate and posterior palate. A number of different systems of nomenclature have been used to demarcate the boundaries of the anterior, medial and posterior regions of the SP. In this thesis, I will follow the nomenclature proposed by Smith et al., (2013) which defines the anterior and posterior palate as tissue lying anterior or posterior to the first molar tooth bud, respectively. The medial palate is defined as palate tissue in the plane of the molar tooth bud.

Several transcription factor genes are differentially expressed along the A-P axis of the developing palatal shelves. The development and maintenance of the expression of these genes in a region specific manner is critical for normal SP development. *Msx1*, *Bmp4*, *Bmp2*, *Shh*, *Spry2*, *Fgf10*, *Fgf7*, *Shox2* genes are preferentially expressed in the anterior palatal mesenchyme (Alappat et al., 2005; Rice et al., 2004; Yu et al., 2005; Zhang et al., 2002), whereas *Barx1*, *Mn1*, *Meox2*, *Tbx22* genes are preferentially expressed in the posterior palatal mesenchyme (Barlow et al., 1999; Li and Ding, 2007; Liu et al., 2008). Although mice lacking either *Msx1* or *Mn1* exhibited complete cleft palate, *Msx1* null mice exhibited specific cell proliferation defects in the anterior region, whereas *Mn1* null mice showed growth deficits in only the middle and posterior regions of the palatal shelves (Zhang et al., 2002; Liu et al., 2008). Mice lacking *Shox2* exhibited a cleft within the anterior palate, whereas the posterior palate fused normally, demonstrating a specific requirement for *Shox2* in anterior palatal outgrowth (Yu et al., 2005). *Tbx22* null mice displayed cleft palate, with the severity varying from a complete cleft palate phenotype due to decreased palatal shelf extension to submucous cleft palate (abnormal muscle orientation in the soft palate during midline fusion) in which palatal shelf elevation and fusion occurred normally (Pauws et al., 2009). The expression of *Msx1* and *Shox2* in the anterior palatal mesenchyme depends on *Bmp* signalling, as expression of both these genes was significantly reduced in the anterior palate of *Wnt1-Cre; Bmpr1af/-* mice (Li et al., 2011).

Most studies on palatal outgrowth have focused on the downward growth of the palatal shelves; however, the maxillary processes also undergo significant rostrocaudal expansion from E12.5 to E14.5, and the palatal shelves correspondingly elongate along their A-P axis. As the palatal shelves elongate,

the domain of *Shox2* mRNA expression expands much more dramatically than that of *Meox2* (Li and Ding, 2007). It has been demonstrated that the A-P boundary of *Shox2* and *Meox2* expression coincides with the first formed palatal rugae (Pantalacci et al., 2008; Welsh and O'Brien, 2009). The rugae are metameric epithelial thickenings on the oral surface of the palate that are first apparent at E12.0 and develop by periodic interposition as the palatal shelves elongate along the A-P axis (Pantalacci et al., 2008). Whereas the second and third rugae form sequentially and anteriorly to the first ruga, the fourth ruga forms between rugae 1 and 2, and, as the palatal shelves continue to elongate along the A-P axis, subsequent rugae formation only occurs between the newly formed ruga and ruga 1 (Pantalacci et al., 2008; Welsh and O'Brien, 2009). Depending on the strain background, mice develop a total of nine or ten rugae, which are maintained after birth.

2.5 Molecular signalling along the O-N axis of the developing SP

The developing palatal shelves also exhibit molecular heterogeneity along the O-N axis during vertical outgrowth with the lateral side forming the oral palatal mesenchyme and the medial side forming the nasal palatal mesenchyme following palatal shelf elevation. Palatal rugae begin to form on the oral side of the developing palatal shelves at E12.0, and the expression of *Shh* becomes restricted to the oral palatal epithelium. *Osr1* and *Osr2* exhibit graded expression in the developing palatal mesenchyme along the oral-nasal axis (Lan et al., 2004). At E13.5, *Osr1* mRNA expression is restricted to the oral side, whereas *Osr2* exhibits graded expression that is strongest in the oral mesenchyme and weaker in the nasal mesenchyme (Yu et al., 2013). Whereas *Pax9* is expressed throughout the O-N axis of the palatal mesenchyme in the posterior region, its expression has been reported to be lower in the oral half

than in the nasal half of the palatal mesenchyme in the anterior region (Yu et al., 2013). *Dlx5* has also been implicated in O-N patterning and palatal expansion. *Dlx5* is co-expressed with *Fgf7* in the oral half of the palatal mesenchyme, and *Fgf7* expression was found to be dramatically reduced in this domain in *Dlx5* null palatal shelves (Han et al., 2009). Although *Dlx5* null palatal shelves elevated and fused, the oral aspect of the palate was dramatically expanded and a malformed soft palate was observed (Han et al., 2009).

2.6 The *Six* family of genes

The *Six* gene family consists of vertebrate genes that have sequence similarities to the *Drosophila* homeobox gene, *Sine oculis* (*So*). These genes are characterized by the presence of a divergent DNA-binding homeodomain (HD, 60 amino acids) and an upstream region, the *Six* domain (SD, 110-115 amino acids), that could be involved in both determining DNA binding specificity and in mediating protein-protein interactions (Fig. 7). In mammals, several members of the family have been cloned and designated *Six1* to *Six6* (Boucher et al., 1995,1996; Oliver et al., 1995; Gallardo et al., 1999; Granadino et al., 1999; Kawakami et al., 1996). The *Six* and homeodomain regions are highly conserved across the *Six* family, while other regions of these proteins are often conserved across vertebrate species, but not between family members. Mouse *Six1* and *Six2* exhibit a high degree of sequence similarity to each other (95% amino acid sequence identity over a region spanning the *Six* domain and homeodomains) and they also show a higher degree of sequence similarity to *Drosophila* *Sine oculis* (*So*) (87 and 95%, respectively, over the same region) than do other family members. *Six1* and *Six2* thus represent orthologues of *Sine oculis* (Oliver et al., 1995). Their expression patterns, however, differ from

that of *So*, which is primarily expressed in the developing visual system of the fly. In mice, fetal expression studies have demonstrated that neither *Six1* nor *Six2* is expressed in the developing eye, rather, there is notable expression in anterior neural structures, neural crest derived tissues, and head, trunk and limb mesenchyme, with *Six1* and *Six2* transcripts appearing to occupy complementary axial domains in the phalanges (Oliver et al., 1995; Ohto et al., 1998).

2.7 Expression patterns of *Six* genes

Six genes are diversely expressed during vertebrate embryogenesis and control the formation of various organs. In most cases, they positively regulate cell numbers within the developing embryo by promoting cell proliferation and survival and mainly act as transcription factors (Christensen et al., 2008; Kawakami et al., 2000).

In the mouse embryo, *Six3* and *Six6* are expressed in restricted areas: developing eye, brain and a few related sites. The expression patterns of these two mouse *Six* genes appear to share similar characteristics with *Drosophila Optix*, whose expression is detected only in the cephalic region of the embryo. In contrast, *Six1*, *Six2* and *Six4* show much broader expression. *Six1* and *Six4* show remarkably similar patterns of expression. They are expressed in two of the sensory organs in the head region (otic vesicles and nasal pits) and in the branchial arches. In addition, the expression of *Six1* and *Six4* extends posteriorly in dorsal root ganglia and somites which give rise to the skeletal muscle of the trunk and limbs. The expression in somites is similar to mesodermal expression of *So* and *D-Six4* in *Drosophila*. *Six2* is also expressed caudally in certain tissues, in addition to the rostral and cervical regions. For *Six5*, the expression pattern was studied by using transgenic mice carrying a

transgene containing its 4.3 kb promoter fragment fused to the *lacZ* reporter gene (Fig. 8). The mice showed expression of the reporter *lacZ* in neural, sensory, muscle, cartilage and other tissues (Heath et al., 1997).



Figure 7: A schematic illustration of the human *SIX2* gene structure showing transcribed exonic sequences present in cloned cDNA (boxes) and the relative locations of the sequences encoding the SIX-(SIX) and homeodomains (HD). (Reproduced with permission from Boucher, C., Carey, N., Edwards, Y., Siciliano, M. and Johnson, K. (1996). Cloning of the human *SIX1* gene and its assignment to chromosome 14. *Genomics* 33: 140-142., via Elsevier and copyright clearance centre).




Figure 8: A schematic drawing of embryonic Expression patterns and chromosomal locations of mouse *Six* family genes. Similarity in the colour of shadings reflects similarity in the expression patterns. DRG, dorsal root ganglion; FB, forebrain; FL, forelimb; GE, genital eminence; HL, hindlimb; NC, nephrogenic cord; NP, nasal placode; OC, optic chiasm; OP, optic vesicle; OV, otic vesicle; RP, Rathke's pouch; SO, somites. B: Expression of *Six3* and *Six6* in a developing eye (E13.5). LE, lens; NR, neural retina; RPE, retinal pigmented epithelium. C: Chromosomal localization of mouse *Six* family genes.

Reproduced with permission from Kawakami, K., Sato, S., Ozaki, H. and Ikeda, K. (2000). *Six* family genes—structure and function as transcription factors and their roles in development. *BioEssays* 22: 616-626., via John Wiley and Sons and Copyright clearance centre.

2.8 *SIX* genes and genetic diseases

Holoprosencephaly (HPE) is a common malformation found in spontaneously aborted fetuses (1/250) and more rarely in live births (1/16,000). It is characterized by signs of abnormal development such as malformation of the brain, a single central incisor, hypotelorism, microcephaly and other craniofacial abnormalities (Roessler et al., 1998).

When the mouse *Six3* gene was mapped to the distal region of chromosome 17, (Oliver et al., 1995) which is orthologous to human 2p21 where the HPE2 locus resides, the possibility that the *SIX3* gene is involved in HPE was suggested. This idea was confirmed by the isolation of the *SIX3* gene from the HPE2 minimal critical region and the identification of four point mutations associated with HPE2 in the coding region of *SIX3*. All of those mutations were identified in the HD of *SIX3*; L226V and del(N232-S234) change amino acid residues unique to the *SIX3/6* subfamily while V250A and R257P change residues common to all Six proteins (Wallis et al., 1999). This strongly suggested that *SIX3* is the HPE2 gene essential for the development of the anterior neural plate and eye in humans.

Human *SIX6* haploinsufficiency was found to be responsible for bilateral anophthalmia, the absence of the optic nerve and chiasma, and for pituitary abnormalities (Gallardo et al., 1999). One such patient was found to have interstitial deletions of the 14q22.3-q23 chromosomal region. The regulatory region and the coding region of *SIX6* were both deleted in one chromosome 14 of that patient. The relationships between *SIX3/6* subfamily genes and several other genes implicated in HPE, such as *SHH*, (Belloni et al., 1996) *Zic2*, (Brown et al., 1998) and TG-interacting factor (*TGIF*) (Walsh et al., 1999) have not yet been clarified.

Myotonic dystrophy (DM) is characterized by muscle weakness, wasting and disturbances in the nervous system (Harris et al., 1996). In addition, DM is associated with ophthalmic features such as cataract, as well as cardiac conduction defects, mental impairment, premature balding and reduced fertility. *SIX5* was originally identified as DMAHP (DM associated homeodomain protein gene), which resides downstream of the causative CTG repeat located on 19q13.3 (DM1) (Boucher et al., 1995). Expansion of the CTG repeat results in increased severity of symptoms and an earlier age of onset. The CTG expansion at this locus has been proposed to alter the local chromatin structure and to repress the transcription of surrounding genes (Otten et al., 1995).

The expression of *SIX5* was reduced 2- to 4-fold in cells derived from DM1 patients relative to wild-type controls (Klesert et al., 1997; Thornton et al., 1997). At least some symptoms were believed to be caused by down-regulation of *SIX5*, which in turn leads to aberrant regulation of the *SIX5* target genes. For example, the observation that myogenin expression is up-regulated most efficiently by *Six5* and *Eya3* (one of the *Eya* family proteins in mouse) through the MEF3 site in the proximal promoter suggests the involvement of *SIX5* in muscle phenotype. The expression of *SIX5* but not DMPK in human lens also suggests the involvement of *SIX5* in the formation of cataracts (Winchester et al., 1999). The involvement of DMPK (DM protein kinase) and DMWD (DM protein with WD repeat), genes located 50 bp downstream to the CTG repeat, has also been suggested which would indicate that DM1 is a multi-gene disorder (Winchester et al., 1999; Alwazzan et al., 1999).

2.9 The *Six2* gene

Six2 is a homeobox transcription factor that is expressed in multiple organ systems of the developing embryo. This gene has been shown to be involved in regulating craniofacial and kidney development (Fogelgren et al., 2008; Fogelgren et al., 2009; Lozanoff, 1993; Lozanoff et al., 2001; Ma and Lozanoff, 1996; Ma and Lozanoff, 1999; McBratney et al., 2003; Singh et al., 1998). In addition, *Six2* has been reported to be expressed in the nasal prominences of E11.5 mice (Brodbeck et al., 2004; Fogelgren et al., 2008; Ohto et al., 1998; Oliver et al., 1995). *Six2* has been shown to be a downstream target of the Hox11 proteins in metanephric kidney development. In the early metanephric mesenchyme, Hox11 proteins interact with *Pax2* and *Eya1* to promote activation of *Six2* (Gong et al., 2007; Wellik et al., 2002). It has also been reported that *Six2* null mice die at birth and exhibit a shorter cranial base. In addition, these mice were also found to have defects in third and fourth branchial arch derived cartilages (Guiyuan et al., 2010). *Six2* has been reported to positively regulate cell numbers within the developing embryo by promoting cell proliferation and survival (Christensen et al., 2008; Kawakami et al., 2000). *Six2* has been implicated in kidney development where it maintains the renal progenitor cell population in an undifferentiated state (Self et al., 2006).

The epigenetic regulation of *Six2* by micro RNAs has also been reported. Lyu et al., (2013) reported that miR-181b directly targets the 3'UTR of *Six2* to downregulate its expression at both the mRNA and protein level in the kidney mesenchyme. In addition, Xiaoyan et al., (2014) have also reported that a single micro RNA (miRNA), miR181c was able to down regulate the expression of *Six2* in the kidney mesenchyme and thus restraining cell proliferation and promoting apoptosis. Furthermore, *Six2* has been reported to

promote breast cancer metastasis through the transcriptional and epigenetic control of *E-cadherin* expression (Wang et al., 2014).

Six2 is a direct downstream target of *Hoxa2*, a gene that plays a direct role in murine secondary palate development (Smith et al., 2009). Expression of *Six2* in wild type embryos is restricted to the first branchial arch mesenchyme. Loss of function of *Hoxa2* results in an expansion of *Six2* expression in the branchial arches and periodic mesenchyme and leads to malformations of the middle and external ear (Kujetova et al., 2005; Rijli et al., 1993). This implies that *Hoxa2* expression is sufficient to repress the expression of *Six2* (Santagati et al., 2005). Interestingly, when *Six2* is overexpressed in the areas of the embryo controlled by *Hoxa2*, the result is a phenotype similar to the *Hoxa2* mutant phenotype. Analysis of the promoter region of *Six2* in combination with ChIP experiments suggests that *Hoxa2* directly represses the expression of *Six2* (Kutejova et al., 2005; Kutejova et al., 2008).

The studies above show the extent to which the *Six2* gene has been studied in mammalian development. Most of these studies have focussed on kidney development and craniofacial development in general. However, no detailed expression analysis of *Six2* has been performed in the developing secondary palate. In addition, its precise role in the craniofacial region and especially in the developing mammalian secondary palate is still unknown. My thesis therefore seeks to address these gaps in knowledge regarding the transcription factor *Six2* during murine palatogenesis.

3. HYPOTHESES

1. Development of the secondary palate is accompanied by changes in the temporal and spatial expression patterns of *Six2* gene transcripts and protein.
2. The *Six2* gene and protein are up-regulated in the absence of the *Hoxa2* gene in the secondary palate.
3. The *Six2* gene regulates cell proliferation during secondary palate development.

4. OBJECTIVES

1. To study the temporal and spatial expression pattern of the *Six2* gene and protein in the developing secondary palate.
2. To investigate whether *Six2* mRNA and protein are altered in the absence of the *Hoxa2* gene in the secondary palate
3. To determine if the *Six2* gene regulates cell proliferation during secondary palate development.

5. MATERIALS AND METHODS

5.1 Hoxa2 transgenic mice

Hoxa2^{+/+}, *Hoxa2*^{+/-} and *Hoxa2*^{-/-} C57 black mice (Gendron-Maguire et al., 1993) were obtained by timed heterozygous matings. Heterozygous animals were obtained by backcrossing heterozygous males with wild-type females. After pups were weaned, animals were anaesthetized using isoflurane 3% in oxygen and approximately 0.2 cm piece of the tail was docked. Genotypes were confirmed by polymerase chain reaction (PCR) analyses (Appendix Fig. 2). Timed heterozygous matings were then performed, and were staged according to Kaufman with mice being considered E0 days pregnant on the day the vaginal plug was found (Kaufman, 1992).

5.2 PCR analysis of genotype

Tissue sample (tail or embryonic tissue) was digested overnight at 55°C in a solution containing 0.5M EDTA, Nuclei lysis solution (Promega) and 20 mg/ml proteinase K enzyme. Following the digestion, samples were allowed to cool to RTM after which protein precipitation solution (Promega) was added and the solution vortexed vigorously at high speed for 20 sec. Solution was then chilled on ice for 5 min followed by centrifugation for 4 min at 14,000 x g. The precipitated protein formed a tight white pellet. The supernatant containing the DNA was then carefully removed (leaving the protein pellet behind) and transferred into a clean 1.5ml microcentrifuge tube containing an equal volume of isopropanol at RTM. The solutions were then mixed gently by inversion until the white thread-like strands of DNA formed a visible mass. The solutions were centrifuged for 1 min at 14,000 x g at RTM. The supernatant

was carefully decanted and the DNA was visible as a small white pellet. Ethanol (70%) at RTM was then added and the tubes gently inverted several times to wash the DNA. The solutions were centrifuged again for 1 min at 14,000 x g at RTM. Ethanol was then carefully aspirated using a Pasteur pipette and the pellet air dried for 10-15 min. The DNA was then rehydrated by adding DNA rehydration solution (Promega) and incubating at 65°C for 1 h. DNA was then analyzed by PCR to determine the genotype of the animal. Each sample was amplified in two different PCR reactions, one to determine the presence of the wild-type *Hoxa2* allele (5'-GTTGGAAGTACCTCTCTTG-3' and 5'-GGGTCCGAGCAGGGTTATTCC-3') and the other to detect the presence of the neomycin cassette that replaces part of exon 1, all of intron 1 and part of exon 2 in the knockout animals (5'-TCGCCTTCTATCGCCTTCTTG-3' and 5'-GTTGGTGTACGCGTTCTCAG-3'). PCR reactions (50 µl) containing 10X amplification buffer (Invitrogen), 2 mM dNTP mix (Sigma), 1µM of each primer (Invitrogen) and 2.5 U of Taq polymerase enzyme (VWR) and 1 µg of DNA were prepared. Reactions were run on the BioRad MyCycler™ using the temperature cycling conditions 95°C for 4 min followed by 35 cycles of 95°C for 30 s, 65°C for 3 min and 72°C for 7 min (Gendron-Maguire et al., 1993). PCR samples were then separated on a 1% agarose gel containing Redsafe™ (Frogga Bio) and visualized using UV light.

5.3 Immunohistochemistry (IHC)

Embryos were harvested from timed-pregnant mice, and heads were fixed in 4% paraformaldehyde in phosphate buffered saline (PBS, pH7.4) for 24 h. Embryo heads were then placed in 30% sucrose in PBS for at least 24 h (or after sinking to bottom of well) before being embedded in optimal cutting temperature (OCT) compound (Tissue-Tek) and serially sectioned (in a coronal

plane) on to 0.5% gelatin coated glass slides. Determination of anterior, medial and posterior was performed by matching to sections from each region (Rice et al., 2004; Smith et al., 2013). Sections were then allowed to dry at RTM for at least 2 h. The 8 μ m thick sections were then rehydrated for 45 min in 1xPBS and blocked in 4% skim milk powder and 0.1% Triton X-100 in PBS for 30 min at RTM. Sections were then exposed to the primary antibody diluted in PBS overnight at 4°C using the following antibodies; *Six2* rabbit polyclonal antibody (ProteintechTM; 1:500), E-cadherin rat monoclonal antibody (SigmaTM; 1:200) and *Ki-67* rat monoclonal antibody (Affymetrix eBioscience; 1:200). Excess unbound primary antibody was rinsed off by two 5 min washes in PBS before being exposed to a labelled secondary antibody for 1.5 h at RTM (IgG Alexa fluor 488 antibody, 1:200 or IgG Alexa fluor 594, 1:400; InvitrogenTM). Finally, sections were rinsed two more times in PBS, mounted in ProLong[®] Gold antifade reagent with DAPI (Molecular Probes) and viewed by fluorescence microscopy. To ensure that the signal observed was not the result of non-specific signal from the secondary antibody, negative controls were performed. These included staining in the absence of the primary antibody, and another set where the secondary antibody was omitted.

5.4 RNA Isolation and Reverse Transcription

Total RNA was isolated from excised wild-type and *Hoxa2* null embryo palate shelves using the RNeasy[®] Protect Mini Kit (Qiagen) as per the manufacturer's protocol. RNA concentration was determined by optical density (Nanovue, General Electric). First strand DNA synthesis was performed using the SuperScript[®] first-strand synthesis system for RT-PCR (Invitrogen), using random decamer primers, as per the manufacturer's protocol. Final concentration of RNA for all RT reactions was 20 ng/ μ l.

5.5 Quantitative Real Time RT-PCR

Gene expression analysis was performed as previously described by Smith et al., 2009. Briefly, Taqman® primers and labelled probe systems (Applied Biosystems) and a real time PCR machine (ABI 7300) from Applied Biosystems were used for real time PCR analysis of *Six2*, *Msx1*, *Bmp4*, *Barx1* and *Ptx1*. Palatal shelves were aseptically dissected from both wild-type and *Hoxa2* null mice at stages E12.5, E13.5, E14.5 and E15.5. All reactions were performed using the Taqman Universal Master Mix (2X), FAM-labelled Taqman Gene Expression assays for *Six2*, *Msx1*, *Bmp4*, *Barx1* and *Ptx1*, VIC-labelled Taqman Endogenous Control β -Actin, and 10ng of cDNA. All reactions were run in replicates of 5 (5 samples, each from a different embryo). Thermocycling parameters were as follows: 2 min at 50°C, 10 min at 95 °C, 40 cycles of 15s at 95 °C followed at end with one cycle at 60 °C for 70s. Taqman gene expression assays have all been tested to have efficiencies not significantly different from 1 (Applied Biosystems, 2004). However, in order to ensure that all primer sets were working properly and that multiplexing the gene specific and β -Actin primers did not result in altered amplification efficiencies all primer sets were examined individually and in complex with β -Actin.

5.6 Droplet digital PCR (ddPCR)

The procedure for PCR analysis using droplet digital PCR was performed as described by Hindson et al., 2013. Briefly, each 20 μ l PCR reaction was loaded into an 8-channel, single-use consumable droplet generation cartridge. Oil (60 μ l) containing emulsion-stabilizing, biocompatible surfactant was loaded into adjacent oil wells and the microfluidic chip was loaded into a beta-series prototype droplet generator (DG). The DG applies a vacuum to the

outlet well creating a pressure difference across the cartridge that simultaneously partitions the sample present in each of the 8 wells into approximately 20,000 monodisperse droplets of accurately known volume. This process occurs at a rate of approximately 1,000 droplets/s per well. The resulting water-in-oil emulsions were pipette-transferred from the outlet well to a 96-well polypropylene plate (Eppendorf), sealed with foil and then amplified to endpoint using a Tetrad2 Peltier Thermal Cycler (Bio-Rad) and the cycling protocol: 95 °C for 10 min then 40 cycles of 95 °C for 15 s and 60 °C for 1 min (2.5 °C/s ramp rate) with a final 10 min hold at 98 °C. Plates containing amplified droplets were loaded into a QX 100 droplet reader (Bio-Rad), which aspirates droplets from the 96-well plate, one well at a time, and streams them single file (about 1,500 droplets/s) past a two-color detector of FAM and VIC fluorescent dyes (Life Technologies) sampling at 100 kHz. Discrimination between droplets that did not contain target (negatives) and those that did (positives) was achieved by applying a global fluorescence amplitude threshold. Concentration estimates were based on the fraction of droplets where amplification has occurred modelled as a Poisson distribution.

5.7 Western blot analysis

Palatal shelves were aseptically dissected from both wild-type and *Hoxa2* null mice before being pooled together prior to lysis in radio-immuno precipitation assay [RIPA] buffer containing protease inhibitor (150 mM NaCl, 10 mM Tris, 0.1%-SDS, 1% Triton X-100, 1% Deoxycholate, 5 mM EDTA). Protein was quantified using the Bradford assay to ensure that equal loading was achieved. Samples were then boiled for 15 min with loading buffer and loaded on a 10% polyacrylamide-SDS gel. After gel separation, the proteins

were transferred to a PolyScreen PVDF membrane, which was subsequently blocked by overnight incubation in 5% skim milk in PBS (SM-PBS) at 4°C.

The membrane was then exposed to a *Six2* primary antibody (*Six2* rabbit polyclonal, Protein tech, 1:2000 in 5% skim milk in PBS) and incubated for 1 h at RTM. This was followed by 4 consecutive washes of 15 min each in PBS with 0.08% Tween-20. After the washes were complete, the membrane was incubated with a secondary antibody specific to the species of the primary antibody that is used (anti-rabbit IgG horseradish peroxidase conjugate (Bio-Rad) 1:3000) in SM-PBS for 1 h at RTM. After 4 washes of 15 min each in PBS-Tween-20, membrane was exposed with a chemiluminescence reagent (ECL substrate, BIORAD) and the signal was detected using the SYNGENE imager. The membrane was then washed overnight at 4°C in PBS before being incubated with the anti- β -tubulin (Development Studies Hybridoma Bank, U. Iowa) at a dilution of 1:2000, and incubated for 1 h at RTM, followed by 4 washes of 15 min each in PBS with 0.08% Tween-20. After the washes, the membrane was exposed to the anti-mouse IgM horseradish peroxidase conjugate (Bio-Rad) in SM-PBS at a dilution of 1:1500, and then treated as indicated above.

After imaging, semi-quantitative densitometry was performed using the Alphamager® densitometric software to determine an integrated density value (IDV) for each band. Four separate western blot assays, each having wild-type and *Hoxa2* null samples of E12.5, E13.5, E14.5, E15.5 were performed. The software allowed for the selection of a representative background area and then to define an area around the bands of interest. Equal size boxes were used in all comparisons. To compare expression, relative IDV values for each blot were calculated by normalizing to β -tubulin levels from the same sample. The resulting values (wild-type and *Hoxa2* null of E12.5, E13.5, E14.5, E15.5)

were then normalized to the E12.5 wild-type sample of that blot. Due to differences in the IDV values between different blots it was necessary to normalize each blot to a band on its own gel to accurately represent any changes in protein expression observed. A total of four blots (n = 4) were analyzed using two way ANOVA to compare Six2 protein expression in wild-type and *Hoxa2* null samples at (E12.5, E13.5, E14.5 and E15.5).

5.8 Culturing MEPM cells

Mouse embryonic palate mesenchyme (MEPM) cells were cultured as previously described by Iwata et al., 2012. Briefly, timed pregnant C57 black mice were sacrificed at stage E13.5 and their embryos removed. Palatal shelves were aseptically excised from the embryo and placed in Hank's balanced salt solution. A piece of tissue was also taken from the embryo for PCR genotyping analysis. In addition, the palatal shelves were treated according to Tyler et al., 1980, to separate the palatal mesenchyme from the epithelium. Purified palatal mesenchyme was incubated with 0.25% trypsin/0.02% EDTA in Ca^{++} - Mg^{++} free PBS for 20 min at 37°C to dissociate individual cells. The digested samples were then briefly triturated and filtered through a 70 μm mesh. The action of trypsin was terminated using 4°C Dulbecco's Modified Eagle's Medium (DMEM)/Ham's medium F12 (1:1) containing 10% fetal bovine serum (FBS) supplemented with 1% penicillin/streptomycin, L-glutamate, sodium pyruvate and non-essential amino acids. The cell number was determined using the MOXI Z automated cell counter (Orflo Technologies) and the primary culture of MEPM cells was initiated by seeding 5×10^5 cells/5ml of complete DMEM/F-12 media in a 25ml tissue culture flask. Cells were sub cultured at a ratio of 1:3 after achieving confluence in a 95% air and 5% CO_2 atmosphere.

5.9 siRNA treatment and cell proliferation analysis

The siRNA used was the pre-designed *Six2* Silencer Select siRNA (Invitrogen: catalogue number 4390771), together with a Silencer select negative control siRNA (Invitrogen). siRNA aliquots were prepared by diluting the stock siRNA (5nM lyophilized powder) using nuclease-free sterile water (Invitrogen) to achieve a working concentration of 20 μ M. MEPM cells (5.0×10^3) were plated in a 96-well plate until they reached 60%-80% confluence. The siRNA sample was mixed with lipofectamine 3000 (Invitrogen) according to the manufacturer's protocol. Briefly, for each transfection, 0.3 μ l of lipofectamine 3000 (Invitrogen) was diluted with 5 μ l of serum free DMEM/F12 media and kept at RTM for 5 min. This was mixed with 5 μ l of serum free media containing 0.25 μ l of siRNA and the mixture vortexed for 10 sec and left at RTM for 20 min. Prior to transfection, cells were rinsed twice with serum free media. The above siRNA-lipofectamine complex (10 μ l) was added to each well of MEPM cells containing 100 μ l of serum free media. Following incubation at 37°C for 12 h, the transfection media was replaced with fresh complete media.

MEPM cells were analyzed for cellular DNA content 48 h from the time of transfection using the CyQUANT NF cell proliferation assay kit (Life Technologies). The cell proliferation assay was performed as per the manufacturer's protocol. Briefly, the transfected MEPM cells were incubated with 1X dye binding solution at 37°C for 45 min in the dark. Fluorescence was detected with a microplate reader (BioTek) with excitation at 485nm and emission at 530nm.

5.10 Statistical analysis

All statistical analyses and graphs were performed using the GraphPad 5.0 software package (GraphPad Prism). Statistical analysis on quantitative real time RT-PCR data for *Six2* expression was evaluated using two-way analysis of variance (2 way ANOVA) followed by Bonferroni post-hoc tests to compare between the genotypes and embryonic stages.

6. RESULTS

6.1 *Six2* protein exhibits a temporal distribution pattern in the developing secondary palate

In order to investigate if *Six2* might be involved in palatogenesis, I first examined whether this gene was endogenously expressed during the different stages of palatogenesis: emergence and vertical outgrowth of palatal shelves (E12.5-E13.5); palatal shelf elevation and horizontal growth (E14.0-E14.5); and palatal shelf fusion (E15.0-E15.5). I utilized fluorescence immunohistochemistry (IHC) to demonstrate that *Six2* protein is expressed differentially across the four stages of secondary palate development (E12.5, E13.5, E14.5, E15.5). Immunohistochemical analysis of coronal sections from the anterior, medial and posterior regions of wild-type embryos showed that *Six2* protein was initially high at E12.5, increased to peak expression at E13.5 and reduced at E14.5 before further declining at E15.5 (Fig. 9).




Figure 9: Six2 protein exhibits a temporal distribution pattern in the developing palate. Immunohistochemical staining of coronal sections from the anterior, medial and posterior regions of wild type embryos showing more staining of Six2 protein at earlier stages of palatogenesis (E12.5-E13.5) and less staining at the later stages (E14.5-E15.5). Panels D, H, I, P are control sections where only the secondary antibody was used. Anterior, medial, posterior and control sections are all from the same embryo. Arrow indicates epithelium, T-tongue, M-mesenchyme, 10X magnification. (scale bar, 10 μ m).

6.2 Six2 protein is spatially distributed in both the epithelium and mesenchyme of the developing secondary palate

Immunohistochemical analyses of the anterior, medial and posterior sections from both wild-type and *Hoxa2* knock out embryos revealed that Six2 protein was consistently expressed in both the epithelium and mesenchyme of mouse palatal shelves across the stages of palatogenesis. It was interesting to note that Six2 protein was expressed in the nasal palatal epithelium (Fig. 10B) but was completely absent from the oral palatal epithelium (Fig. 10D). This expression pattern was consistent in the A-P axis of the palate and across the stages of secondary palate development. To confirm this finding, I double-labelled palatal sections by IHC using *Six2* and *E-cadherin* antibodies. *E-cadherin* is a known epithelial marker that stains only in the epithelium. Cells in the nasal palatal epithelium were positive for both *Six2* and *E-cadherin* staining (Fig. 10J), while cells in the oral palatal epithelium were only positive for *E-cadherin* (Fig. 10L). This confirmed my earlier observation that Six2 protein was only expressed in the nasal palatal epithelium and was completely absent from the oral palatal epithelium.

Furthermore, I observed more Six2 protein expression in the oral half of the palatal mesenchyme as compared to the nasal half of the palate mesenchyme (Fig. 11). This pattern of expression was again consistent in the A-P axis of the palate and from E12.5 to E14.5. At E15.5 however, Six2 protein expression seemed to be concentrated in the central portions of the palatal mesenchyme when compared to the oral and nasal halves of the palatal mesenchyme where expression was less.




Figure 10: Six2 protein exhibits a spatial expression pattern in the palate epithelium. Immunohistochemical double-labelling of Six2 protein (A,B,C,D) and of E-cadherin (E,F,G,H) on coronal sections of the developing palate at E14.5. Staining of Six2 protein can be observed in the nasal epithelium (B), reduced staining in the region of the midline epithelial seam (MES) (C) and no staining in the oral epithelium (D). Blue boxes in panels A, E represent magnified regions (40X) for the MES, nasal and oral epithelium as shown in panels B,C,D,F,G,H,J,K & L . Sections I,J,K,L are overlays. (Scale bar, 10 μ m).




Figure 11: Six2 protein exhibits a spatial expression pattern in the palatal mesenchyme. Immunohistochemical staining of representative coronal sections of the developing secondary palate of wild-type embryos (E12.5-E15.5). From E12.5 to E14.5, there is more staining for Six2 protein in the oral half of the palatal mesenchyme (arrows) and less staining in the nasal half of the palatal mesenchyme (asterisks) (A-C). At E15.5, Six2 protein expression is concentrated in the central portions of the palatal mesenchyme (arrows) (D). T-tongue.(10x Magnification) (Scale bar, 10 μ m).


6.3 Six2 protein exhibits an A-P differential expression pattern in the developing secondary palate

Analysis of the anterior, medial and posterior regions of the palatal epithelium and mesenchyme revealed a differential expression pattern of Six2 protein in the A-P axis in both the wild-type and *Hoxa2* null embryos. *In vivo* cell counting analysis revealed a general increase in the A-P direction of the number of palate mesenchymal cells expressing Six2 protein, with the least expression in the anterior and most expression in the posterior regions of the palate (Fig. 12A).

In addition, IHC analysis showed increasing Six2 protein expression in the A-P direction with the most expression observed in the posterior portions of the palate mesenchyme (Fig. 12B). This expression pattern was consistent for both wild-type and *Hoxa2* null embryos and across the stages of secondary palate development. Also interestingly, it seems that Six2 protein is primarily expressed in the anterior nasal palatal epithelium and it progressively decreases as one moves into the posterior regions where little or no expression is seen in the posterior nasal palatal epithelia (Fig. 12B).

A)

In vivo cell counting analysis of Six2 positive cells at E13.5



B)




Figure 12: Six2 protein exhibits an Anterior-Posterior differential expression pattern in the palate epithelium and mesenchyme.

A) *In vivo* cell counting analysis of Six2 protein positive cells from coronal anterior, medial and posterior sections of wild-type and *Hoxa2* null mice (E13.5) showing highest Six2 protein expression in the posterior region of the palatal mesenchyme of *Hoxa2* null embryos.


B) Representative coronal sections from the anterior, medial and posterior regions of wild type embryos (E13.5) showing Six2 protein expression the palatal mesenchyme increasing from the anterior towards the posterior region of the secondary palate. It is interesting to observe that in the posterior regions, expression of Six2 in the nasal palatal epithelium is also much lower compared to the anterior and medial epithelia. This pattern of expression is observed across all stages of palatogenesis and in both wild-type and *Hoxa2* null mice. Six2 positive cells were expressed as a percentage of the total cell number (DAPI positive cells). Bars represent mean \pm SEM, $n=4$, * $p<0.05$, ** $p<0.01$. T-tongue. (scale bar, 10 μ m).

6.4 *Six2* protein and mRNA are up-regulated in the absence of *Hoxa2*



The *Hoxa2* gene has been reported to bind to the *Six2* promoter *in vivo* to repress its expression (Kujetova et al., 2008). To confirm these findings, I performed quantitative real time PCR (qPCR) and western blot analysis on palatal shelves from both wild-type and *Hoxa2* null embryos. There was a significant increase in the expression of *Six2* at both the mRNA and protein level in the *Hoxa2* null embryos when compared to their wild-type littermates at all stages of palatogenesis, with peak expression occurring at E13.5 (Figs. 13B, C). The peak expression of *Six2* at E13.5 coincides with the time of palatal shelf vertical outgrowth on either sides of the tongue. In addition, I also performed absolute mRNA quantification by droplet digital PCR (ddPCR) to compare between ddPCR and qPCR (Appendix Fig.1). The results for mRNA quantification using ddPCR and qPCR matched indicating consistency of results using the two different techniques (Fig.13C and Appendix Fig.1).

Furthermore, IHC analyses of anterior, medial and posterior sections of both wild-type and *Hoxa2* null embryos across the stages of palatogenesis revealed more *Six2* protein expression in the *Hoxa2* null embryos compared to their wild type littermates (Fig. 13A). Additionally, *in vivo* cell counting analysis of *Six2* protein positive cells in the nasal half of the palatal mesenchyme in wild-type and *Hoxa2* null littermates revealed an expanded domain of *Six2* protein expression in the *Hoxa2* null palatal shelves. In the wild-type palatal shelves, *Six2* protein expression is seen mainly in the oral half of the palatal mesenchyme with less expression observed in the nasal half of the palatal mesenchyme. In the *Hoxa2* null palatal shelves however, the domain of *Six2* protein expression expands from the oral half of the palatal mesenchyme to include the nasal half of the palatal mesenchyme (Fig. 14).

A)



Six2 western blot analysis



Six2 mRNA qPCR analysis




Figure 13: *Six2* mRNA and protein are up-regulated in the absence *Hoxa2*.

A) The domain of *Six2* protein expression is expanded in the *Hoxa2* null palatal shelves. Immunohistochemical staining of representative medial coronal sections of the secondary palate from wild-type and *Hoxa2* null littermates showing an extended domain of mesenchymal *Six2* protein expression in the *Hoxa2* null palatal shelves (E, F, G, H) compared to wild type palatal shelves (A, B, C, D). In the *Hoxa2* null palatal shelves, the domain of *Six2* expression expands from the oral half of the palatal mesenchyme to include the nasal half of the palatal mesenchyme. Also, the palatal shelves fail to elevate above the tongue at E14.5 and E15.5 in the *Hoxa2* null embryos. T-tongue. (scale bar, 10µm).

B) Western blot analysis of protein obtained from palatal shelves of wild-type and *Hoxa2* null mice. There is enhanced *Six2* protein expression in the absence of *Hoxa2* across all stages of palatogenesis with peak expression occurring at E13.5 in both wild-type and *Hoxa2* null palatal shelves.

C) Relative quantitative expression of *Six2* mRNA during murine palatogenesis in wild-type and *Hoxa2* null mice. In the wild type mice, *Six2* mRNA expression is comparably high at E12.5 and E13.5 and reduces at E14.5 before decreasing further at E15.5. In the *Hoxa2* null mice, expression is enhanced across all stages of palatogenesis. In the absence of *Hoxa2*, *Six2* mRNA is initially high at E12.5 and increases to a peak level at E13.5. This expression dramatically reduces at E14.5 before decreasing further at E15.5. Bars represent mean \pm SEM, n=5, *p<0.05, **p<0.01.

In vivo cell counting analysis of Six2 protein positive cells in the nasal half of the palate mesenchyme.




Figure 14: *In vivo* cell counting analysis of Six2 protein positive cells in the nasal half of the palatal mesenchyme in wild-type and *Hoxa2* null littermates (E14.5) showing an expanded domain of Six2 protein expression in the *Hoxa2* null palatal shelves. In the wild-type palatal shelves, Six2 protein expression is seen mainly in the oral half of the palatal mesenchyme with less expression observed in the nasal half of the palatal mesenchyme. In the *Hoxa2* null palatal shelves however, the domain of Six2 protein expression expands from the oral half of the palatal mesenchyme to include the nasal half of the palatal mesenchyme. Six2 positive cells were expressed as a percentage of the total cell number (DAPI positive cells). Bars represent mean \pm SEM, $n=4$, * $p<0.05$, ** $p<0.01$.

6.5 Analysis of cell proliferation in vivo

To investigate whether *Six2* has a role to play during palatal mesenchyme cell proliferation, I performed IHC analyses on anterior, medial and posterior palatal sections of both wild-type and *Hoxa2* null embryos, using an antibody for the Ki-67 antigen, a known marker of cell proliferation. This involved counting cells in a whole palatal shelf including; the total number of cells (DAPI positive cells), cells expressing the *Six2* protein (*Six2* positive cells), actively proliferating cells (Ki-67 positive cells) and the cells that were actively proliferating and co-expressing the *Six2* protein (cells positive for both *Six2* and ki-67). I chose to investigate this putative role of *Six2* at embryonic stage (E) 13.5 when the palatal shelves start growing vertically downwards on either side of the tongue. Incidentally, this is also the stage where I observed peak expression of the *Six2* gene at both the mRNA and protein level (see Figs. 13A, B, C).

I observed a significant increase in cell proliferation in the *Hoxa2* null palatal shelves as compared to the wild-type palatal shelves in both the A-P axis and the O-N axis of the palatal mesenchyme, with the highest percentage of proliferating cells observed in the nasal half of the posterior palatal mesenchyme (Figs. 15A-G). In addition, the percentage of proliferating cells that were also expressing the *Six2* protein (*Six2*/ki-67 double positive cells) was significantly increased in the *Hoxa2* null palatal shelves when compared to the palatal shelves from wild-type embryos, in both the A-P axis and the O-N axis of the palate mesenchyme. Again, the highest percentage of these double positive cells was observed in the nasal half of the posterior palatal mesenchyme (Figs. 16A-G).




Figure 15: Increased cell proliferation in nasal half of the posterior palatal mesenchyme of *Hoxa2* null palatal shelves.

A) Immunohistochemical staining of coronal sections from the anterior, medial and posterior regions of wild-type and *Hoxa2* null palatal shelves (E13.5) for Ki-67, a marker of cell proliferation. **(A-G)** In the *Hoxa2* null palatal shelves, there is a significant increase in cell proliferation in the both the A-P axis and O-N axis, with the highest percentage of proliferating cells observed in the nasal half of the posterior palatal mesenchyme. Asterisks indicate the nasal half of the palatal mesenchyme. Proliferating cells were expressed as a percentage of the total cell number (DAPI positive cells). (scale bar, 10 μ m). Bars represent mean \pm SEM, n=4, *p<0.05, **p<0.01.




Figure 16: The number of cells co-expressing *Six2* and *Ki-67* is highest in the nasal half of the posterior palatal mesenchyme of *Hoxa2* null palatal shelves. **A)** Immunohistochemical staining of coronal sections from the anterior, medial and posterior regions of wild-type and *Hoxa2* null palatal shelves (E13.5) for both *Six2* and *Ki-67* (double staining). **(A-G)** There is a significant increase in the percentage of proliferating cells that are co-expressing *Six2* protein (*Six2*/*Ki-67* double positive cells) in the *Hoxa2* null palatal shelves in both the A-P and O-N axes of the palatal mesenchyme. The highest percentage of these double positive cells was observed in the in the nasal half of the posterior palatal mesenchyme. Asterisks indicate the nasal half of the palatal mesenchyme. Cells co-expressing *Six2* and *Ki-67* were expressed as a percentage of total *Six2* positive cells (scale bar, 10 μ m). Bars represent mean \pm SEM, $n=4$, * $p<0.05$, ** $p<0.01$.

6.6 Analysis of cell proliferation *in vitro*

In addition to *in vivo* cell proliferation analysis, I also performed analyses of cell proliferation *in vitro*. This involved knocking down the expression of *Six2* mRNA in primary palatal cell cultures using *Six2* siRNA and observing the effect this had on cell proliferation and the expression of the cell cycle regulator, *Cyclin D1*. I used an MEPM cell culture system of both wild-type and *Hoxa2* null cells. These were freshly cultured primary cells prepared from the palate mesenchyme of E13.5 embryos, a stage at which palatal shelves grow vertically downwards on either sides of the tongue musculature.

In order to achieve *Six2* mRNA knock down in MEPM cells, two *Six2* siRNA sequences were tested namely; sequence 1 and sequence 2. *Six2* siRNA sequence 1 achieved 90% knockdown of *Six2* mRNA while *Six2* siRNA sequence 2 only achieved 30% knockdown of *Six2* mRNA (Fig. 17). I therefore chose to use *Six2* siRNA sequence 1 for my subsequent experiments.

In addition, I used the CyQUANT® NF Cell Proliferation Assay Kit (Invitrogen) for *in vitro* analysis of cell proliferation. The CyQUANT® NF assay is based on measurement of cellular DNA content via fluorescent dye binding. Because cellular DNA content is highly regulated, it is closely proportional to cell number. The extent of proliferation was determined by comparing cell counts for siRNA treated MEPM cell cultures and the control cell cultures. The two controls used in my experiment were the mock control, where there was no siRNA treatment and the cell cultures treated with a scramble control siRNA from the same vendor (Invitrogen) that is known not to cause *Six2* mRNA knockdown.

It was interesting to observe that *in vitro* *Six2* siRNA knock down restored cell proliferation in the *Hoxa2* null MEPM cell cultures to wild-type levels (Fig. 18A). In both the wild-type and *Hoxa2* null cell cultures, I observed

that cell proliferation was significantly reduced when they were treated with *Six2* siRNA compared to the mock and control siRNA cell cultures.

Furthermore, my study sought to find out if the knockdown of *Six2* *in vitro* would have an effect on the expression of *Cyclin D1*. Indeed, I found that *in vitro* *Six2* siRNA knock down restored *Cyclin D1* expression in the *Hoxa2* null palatal mesenchyme cell cultures to wild-type levels (Fig. 18B). In both the wild-type and *Hoxa2* null palatal mesenchyme cell cultures, *Cyclin D1* expression was significantly reduced after *Six2* siRNA knock down compared to the mock and control siRNA treated cell cultures.

***Six2* mRNA relative Quantification**





Figure 17: *Six2* siRNA knockdown in MEPM cells. 90% *Six2* mRNA knockdown was achieved using *Six2* siRNA sequence-1 compared to 30% *Six2* mRNA knockdown using *Six2* siRNA sequence-2. *Six2* siRNA sequence-1 was therefore chosen for the subsequent experiments. Mock treated cell cultures are those where there was no siRNA treatment, while the control siRNA cell cultures were treated with a scramble control siRNA. Bars represent mean \pm SEM, n=4, *p<0.05, **p<0.01.

A) Cell proliferation assay



B) *Cyclin D1* mRNA expression




Figure 18: A) *In vitro* Six2 siRNA knock down restored cell proliferation in the *Hoxa2* null palate mesenchymal cell cultures to wild-type levels. In both the wild-type and *Hoxa2* null MEPM cell cultures, proliferation was significantly reduced when they were treated with *Six2* siRNA compared to the mock and control siRNA cultures. **B) *In vitro* Six2 siRNA knock down restored *Cyclin D1* expression in the *Hoxa2* null palate mesenchymal cell cultures to wild-type levels.** In both the wild-type and *Hoxa2* null MEPM cell cultures, *Cyclin D1* expression was significantly reduced when they were treated with *Six2* siRNA compared to the mock and control siRNA cultures. Mock treated cell cultures are those where there was no siRNA treatment, while the control siRNA cell cultures were treated with a scramble control siRNA. Bars represent mean \pm SEM, n=4, *p<0.05, **p<0.01.

During early embryonic development, a number of genes and signalling molecules including, *Msx1*, *Bmp4*, *Fgf10*, *Barx1* and *Ptx1* are expressed in the developing secondary palate (Zhang et al., 2002, Yu et al; 2005, Alappat et al., 2005). In this study, I wanted to determine if *in vitro* *Six2* siRNA knock down in MEPM cell cultures would have an effect on the expression of *Bmp4*, *Msx1*, *Barx1* and *Ptx1*; some of the palatal mesenchyme cell signalling molecules expressed during palatogenesis. *In vitro* *Six2* siRNA knock down did not have an effect on the expression of the mRNA transcripts of these genes (Fig. 19). However, the expression of *Bmp4*, *Barx1* and *Ptx1* mRNAs was significantly enhanced in the *Hoxa2* null palatal mesenchyme cell cultures compared to the wild-type cell cultures. In addition, I observed that *Msx1* expression was significantly reduced in the *Hoxa2* null cell cultures compared to the wild-type cell cultures.




Figure 19: *In vitro* Six2 siRNA knock-down did not have an effect on the expression of *Msx1*, *Bmp4*, *Barx1* and *Ptx1* mRNA transcripts. However, the expression of *Bmp4*, *Barx1* and *Ptx1* remained higher in the mock-treated control *Hoxa2* null MEPM cell cultures compared to the mock-treated control wild-type cell cultures. *Msx1* expression remained significantly reduced in the mock-treated control *Hoxa2* null cell cultures compared to the mock-treated control wild-type cell cultures. Bars represent mean \pm SEM, $n=4$, * $p<0.05$, ** $p<0.01$.

7. DISCUSSION

7.1 *Six2* exhibits a temporal-spatial expression profile in the developing secondary palate

In order to study the function of *Six2* in the developing secondary palate, my study first sought to establish if this gene is expressed in the mouse secondary palate. Indeed, *Six2* is expressed in the developing SP and my study has characterized for the first time the temporal and spatial distribution patterns of *Six2* in the developing secondary palate.

Using real time PCR and western blot analysis, I was able to demonstrate that *Six2* mRNA and protein respectively are expressed in the developing SP from the time of palatal shelf emergence at E12.5 to the time of palatal shelf fusion at E15.5 (Fig. 13B, C). In addition, fluorescence IHC on coronal sections from heads of wild-type mouse embryos revealed *Six2* protein expression at all the stages of SP development (Fig.9). *Six2* expression at both the mRNA and protein level was higher during the earlier stages of palatogenesis (E12.5 and E13.5) compared to the later stages of palatogenesis (E14.5 and E15.5) (Fig. 9 and Fig. 13A-C). *Six2* expression is initially high at E12.5, it then increases to a peak expression at E13.5 before dramatically reducing at E14.5 and further declining by the time of palatal shelf fusion at E15.5. It is interesting to note that the peak expression of *Six2* at E13.5 coincides with the time when the paired palatal shelves start to grow vertically downwards on either sides of the tongue.

These findings have expanded on what earlier studies had reported that *Six2* is expressed in the craniofacial region of developing mouse embryos as early as E11.5 (Oliver et al., 1995; Brodbeck et al., 2004; Fogelgren et al., 2008). In addition, the high expression of *Six2* at the earlier stages of SP development

points to the possibility that this gene might be critical during the stages of palatal shelf emergence (E12.5) and palatal shelf vertical outgrowth (E13.5). The rapid decline of *Six2* expression between E14.0 and E15.5 suggests that this gene may not be as critical during the later stages of palatogenesis.

In addition to demonstrating that there are temporal changes in *Six2* expression levels during palatogenesis, my study also revealed regional differences in the spatial expression pattern of *Six2* in the developing SP. Fluorescence IHC demonstrated that *Six2* protein was present in both the palate epithelium and palatal mesenchyme. Whereas *Six2* protein is highly expressed in the nasal palatal epithelium (Fig. 10B and Fig. 20)), its expression is devoid from the oral palatal epithelium (Fig. 10D and Fig. 20). To confirm this finding, I utilized IHC double labelling of *Six2* and E-cadherin, a known epithelial marker. The cells in the nasal palatal epithelium were positive for both *Six2* and E-cadherin (Fig. 10J), while the cells lining the oral palatal epithelium were only positive for E-cadherin (Fig. 10L).

Interestingly, I observed an opposite pattern of *Six2* protein expression in the palatal mesenchyme. *Six2* protein was highly expressed in the oral half of the palatal mesenchyme, while reduced expression was observed in the nasal half of the palate mesenchyme (Fig. 11). This pattern of expression was consistent across the A-P axis of the palate and from E12.5 to E14.5 (Fig. 11A-C and Fig. 20). At E15.5 however, the pattern of *Six2* protein expression in the mesenchyme seemed to be altered, with expression concentrated in the central portions of the fused palate (Fig. 11D).

Previous studies by different groups have also reported genes that exhibit differential expression patterns in the developing SP. *Pax9*, a gene that has been reported to play an important role in regulating epithelial-mesenchymal interactions during SP development is expressed in both the

mesenchyme and posterior palatal epithelium (Yu et al., 2013). Whereas Pax9 is expressed throughout the O-N axis of the palatal mesenchyme in the posterior region, its expression is lower in the oral half than in the nasal half of the palatal mesenchyme in the anterior region. Although the reason for this complimentary pattern of Six2 protein expression in the palate epithelium and mesenchyme is still unclear, it is possible that Six2 could be involved in the epithelial-mesenchymal cross-talk that drives palatal shelf outgrowth.




Figure 20: A schematic diagram illustrating the spatial expression pattern of Six2 protein in the developing palate at E13.5 and E14.5. Whereas there is expression of Six2 protein in the nasal palatal epithelium (block arrows) and the oral half of the palatal mesenchyme (OM), there is no expression in the oral palatal epithelium (stick arrows) and the nasal half of the palatal mesenchyme (NM), T-tongue.

7.2 Six2 protein exhibits an A-P differential expression pattern in the developing secondary palate.

The SP may be divided into three regions along the A-P axis namely; the anterior, medial and posterior palate. Various nomenclature systems have been used to designate the boundaries of the anterior, medial and posterior regions of the SP. When preparing coronal sections for this study, I followed the nomenclature of Smith et al., (2013) which defines tissue lying anterior to the first molar tooth bud as the anterior palate, tissue lying posterior to the first molar tooth bud as the posterior palate, and tissue in the plane of the first molar as the medial palate. Anterior, medial and posterior histological sections were taken from the same embryo. Using IHC analysis, I was able to demonstrate that Six2 protein exhibits an A-P differential expression profile in the developing SP. Although I observed a substantial amount of Six2 protein expression in the anterior regions of the palate mesenchyme, there appeared to be a gradual increase of Six2 protein expression towards the posterior regions of the palate mesenchyme. Also interestingly, it seems that Six2 protein is primarily expressed in the anterior nasal palatal epithelium and it progressively decreases along the A-P axis, such that little or no expression is seen in the most posterior nasal palatal epithelium (Fig. 12).

Previous studies have linked gene expression along the A-P axis to their functions in secondary palate development. For example, *Msx1* is a gene that is primarily expressed in the anterior portions of the SP, while *Mn1* is expressed mostly in the posterior regions of the palate. Whereas mice lacking either *Msx1* or *Mn1* exhibit a complete cleft palate, *Msx1* null mice exhibit specific cell proliferation defects in the anterior region, while *Mn1* null mice have growth deficits in only the middle and posterior regions of the palatal shelves (Zhang et al., 2002; Lin et al., 2008).

A plausible explanation for the disparity in the patterns of *Six2* protein expression in the epithelium and mesenchyme of the SP along the A-P axis could be that in the epithelium, *Six2* protein executes most of its functions in the anterior portions of the SP while in the palatal mesenchyme, most of its functions are performed in the posterior regions of the secondary palate.


7.3 *Six2* mRNA and protein are up-regulated in the absence of *Hoxa2*

A study by Kujetova et al., (2005) reported *Six2* to be an immediate downstream target of *Hoxa2*, a gene that has been found to play a direct role in mouse SP development (Smith et al., 2009). There have also been reports that the expression of *Six2* in the wild-type embryo is restricted to the first branchial arch mesenchyme, and that the loss of function of *Hoxa2* results in an expansion of the domain of *Six2* expression in the branchial arches and in the mesenchyme and leads to malformations of the middle and external ear. This implies that *Hoxa2* expression is sufficient to repress expression of *Six2* (Rijli et al., 1993; Kujetova et al., 2005). In my study, I sought to confirm if loss of *Hoxa2* function indeed alters the expression of *Six2* in the developing SP. My study utilized samples from palatal shelves of both wild-type and *Hoxa2* null embryos. Using qPCR and western blot analysis, I was able to demonstrate that *Six2* mRNA and protein respectively, were consistently up-regulated in the absence of *Hoxa2*, from the time of initial palatal shelf emergence at E12.5, through to the time of palatal shelf fusion at E15.5 (Fig. 13B, C). In addition, IHC analyses of anterior, medial and posterior palatal sections of both wild-type and *Hoxa2* null embryos across the stages of palatogenesis revealed more *Six2* protein expression in the *Hoxa2* null embryos compared to their wild-type littermates (Fig. 13A).

Furthermore, *in vivo* cell counting analysis of Six2 protein positive cells in the nasal half of the palatal mesenchyme in wild-type and *Hoxa2* null littermates revealed an expanded domain of Six2 protein expression in the *Hoxa2* null palatal shelves. In the wild-type palatal shelves, Six2 protein expression was observed mainly in the oral half of the palatal mesenchyme with less expression observed in the nasal half of the palatal mesenchyme. In the *Hoxa2* null palatal shelves however, the domain of Six2 protein expression expanded from the oral half of the palatal mesenchyme to include the nasal half of the palatal mesenchyme where intense staining for Six2 protein was also observed (Figs. 13A, 14, 21). It is interesting to note that unlike the wild-type palatal shelves, the *Hoxa2* null palatal shelves failed to reorient themselves above the tongue but remained positioned vertically on either sides of the tongue (Fig.13A).

My findings therefore confirm that indeed, *Six2* is an immediate downstream target of *Hoxa2* and that *Hoxa2* expression alone is enough to repress the expression of *Six2*.

A. Wild type



B. *Hoxa2* null




Figure 21: A schematic diagram demonstrating an expanded domain of *Six2* protein expression in the mesenchyme of the *Hoxa2* null palatal shelf at E14.5.

A) In the wild-type palatal shelves, *Six2* protein expression is observed mostly in the oral half of the palatal mesenchyme.

B) In the *Hoxa2* null palatal shelves however, the domain of *Six2* protein expression in the mesenchyme expands to include the nasal half of the palatal mesenchyme. Also, unlike the wild-type palatal shelves, the *Hoxa2* null palatal shelves fail to re-orient themselves above the tongue. NE-nasal palatal epithelium, OE-oral palatal epithelium, NM-nasal half of palatal mesenchyme, OM-oral half of palatal mesenchyme, T-tongue.

7.4 *Six2* has an impact on palate mesenchymal cell proliferation during palatogenesis

7.4.1 There is increased cell proliferation and a higher percentage of cells co-expressing *Six2* and Ki-67 in the nasal half of the posterior palatal mesenchyme of *Hoxa2* null palatal shelves.

Six2 has been reported to be an immediate downstream target of *Hoxa2* and is up-regulated in *Hoxa2* null embryos (Santagati et al., 2005). In addition, a study by Smith et al., (2009) reported altered cell proliferation in *Hoxa2* null palatal shelves. My study therefore sought to investigate if the altered cell proliferation observed in the *Hoxa2* null palatal shelves is due to the up-regulation of *Six2*.

In order to investigate this potential role for *Six2* during palate mesenchymal cell proliferation, I performed IHC analyses on anterior, medial and posterior coronal palatal sections of both wild-type and *Hoxa2 null* embryos, using an antibody for the Ki-67 antigen which is a known marker of proliferating cells. I had initially wanted to use BrDU as opposed to Ki-67 but I encountered technical difficulties in double-labelling *Six2* and BrDU. I performed a cell counting analysis on whole palatal shelves and also the oral and nasal halves of the palatal mesenchyme. The cells counted included the total number of cells (Dapi positive cells), cells expressing *Six2* protein (*Six2* positive cells), actively proliferating cells (Ki-67 positive cells) and cells that were actively proliferating and co-expressing *Six2* protein (cells positive for both *Six2* and Ki-67). I chose to investigate this putative role of *Six2* at stage E13.5 because this is the stage at which *Six2* expression is highest at both the mRNA and protein level. Incidentally, this is also the stage when the palatal shelves start to grow vertically downwards on either sides of the tongue. Cell

counting analysis on whole palatal shelves after Ki-67 staining revealed a significant increase in cell proliferation in the *Hoxa2* null palatal shelves compared to the wild-type palatal shelves in both the A-P and O-N axes of the palatal mesenchyme, with the highest percentage of proliferating cells observed in the nasal half of the posterior palatal mesenchyme of *Hoxa2* null palatal shelves (Fig. 15). This finding is in agreement with that of Smith et al., (2009) who also observed increased cell proliferation in the *Hoxa2* null palatal shelves.

Furthermore, I found that the percentage of actively proliferating cells that were co-expressing Six2 (Six2/Ki-67 double positive cells) was significantly increased in the *Hoxa2* null palatal shelves when compared to palatal shelves from wild-type embryos in both the A-P and O-N axes of the palatal mesenchyme, with the highest percentage of these double positive cells observed in the nasal half of the posterior palatal mesenchyme of *Hoxa2* null palatal shelves (Fig. 16).

As mentioned earlier, I observed Six2 protein expression exclusively in the nasal half of the palatal epithelium. In addition, I demonstrated that *Six2* mRNA and protein are up-regulated in the absence of *Hoxa2* and that Six2 protein expression is highest in the posterior regions of the secondary palate. These observations, coupled with the fact that I noticed increased cell proliferation and the highest percentage of actively proliferating cells that were co-expressing Six2 in the nasal half of the posterior palatal mesenchyme of *Hoxa2* null palatal shelves suggests a potential role for *Six2* during the palate epithelial-mesenchymal crosstalk that drives palate mesenchymal cell proliferation and palatal shelf outgrowth.

7.4.2 *In vitro* Six2 siRNA knockdown restored cell proliferation and Cyclin D1 expression in Hoxa2 null MEPM cell cultures to wild-type levels

In order to further investigate the role of *Six2* in palate mesenchymal cell proliferation, I performed *in vitro* *Six2* siRNA knockdown in MEPM primary cells obtained from palatal shelves of both wild-type and *Hoxa2* null mouse embryos. The palatal shelves were harvested at E13.5 when *Six2* expression is at its peak. I mentioned earlier that my study has confirmed previous reports that there is increased palate mesenchymal cell proliferation in *Hoxa2* null palatal shelves compared to wild-type palatal shelves. In addition, my study reveals enhanced *Six2* expression in the absence of *Hoxa2*. It was therefore interesting to discover that *in vitro* *Six2* siRNA knockdown restored palate mesenchymal cell proliferation in the *Hoxa2* null MEPM cell cultures to wild-type levels (Fig. 18A). This finding points to the possibility that the increased cell proliferation observed after knocking out *Hoxa2*, could in part, be due to enhanced *Six2* expression. Zhang et al., (2002) and Rice et al., (2004) found that exogenous application of *Shh* induced a mitogenic response in palatal explants cultures mediated in part by the cell cycle regulators *Cyclin D1* and *Cyclin D2*. In this study, I found that *in vitro* *Six2* siRNA knockdown also restored *Cyclin D1* expression in the *Hoxa2* null MEPM cell cultures to wild-type levels (Fig. 18B). This implies that the cell cycle regulator *Cyclin D1* is acting downstream of *Six2* in the palate mesenchyme. This finding re-affirmed my initial suspicions that *Six2* might play a crucial role in palate mesenchymal cell proliferation.

My study further investigated if *in vitro* *Six2* siRNA knock down had an effect on the expression of *Msx1*, *Bmp4*, *Barx1* and *Ptx1*, some of the genes involved in palate mesenchymal cell signalling during SP development (Smith

et al., 2009). *In vitro* *Six2* siRNA knockdown however, did not alter the expression of any of the above mentioned genes (Fig. 19). A possible explanation for this observation could be that these genes are acting upstream of *Six2* in the mesenchyme during SP development. Collectively, the ability of actively proliferating palate mesenchymal cells to co-express *Six2* protein and the restoration of palate mesenchymal cell proliferation and *Cyclin D1* expression in *Hoxa2* null MEPM cell cultures to wild-type levels points to the possibility that *Six2* might have a role to play in palate mesenchymal cell proliferation. In addition, this finding has confirmed the cell cycle regulator *Cyclin D1* as a putative downstream target of *Six2* in the palate mesenchyme.

8. CONCLUSION

Although *Six2* has been widely studied in mammalian embryonic development, its precise role during mammalian secondary palate development has never been reported. My study reveals for the first time that *Six2* is expressed during all stages of secondary palate development with peak expression occurring at E13.5 at both the mRNA and protein level. My results also show that *Six2* protein is expressed in the nasal palatal epithelium but is completely absent from the oral palatal epithelium. In addition, I observed increased staining for *Six2* protein in the oral half of the palatal mesenchyme and reduced staining for *Six2* protein in the nasal half of the palatal mesenchyme. My study also demonstrates that *Six2* protein exhibits an A-P differential expression pattern in the developing SP. *Six2* protein expression in the palatal mesenchyme increases in the anterior to posterior direction with the most expression observed in the posterior regions of the palate. On the contrary, *Six2* protein expression in the nasal palatal epithelium seems to decrease as one moves from the anterior regions of the palate to the posterior regions of the palate.

Furthermore, I show that *Six2* is up-regulated in the absence of *Hoxa2* at both the mRNA and protein level, consistent with earlier reports that *Hoxa2* binds to the *Six2* promoter *in vivo* to repress its expression. I also successfully demonstrated that actively proliferating palate mesenchymal cells were co-expressing *Six2* protein, indicating a potential role for *Six2* during palate mesenchymal cell proliferation. This finding was further strengthened by *in vitro* studies on primary MEPM cell cultures which showed that *in vitro* *Six2* siRNA knockdown restored cell proliferation and *Cyclin D1* expression in the *Hoxa2* null MEPM cell cultures to wild type levels.

Collectively, my data reveals a novel expression profile for *Six2* in the developing murine SP and suggests a possible role it might play during the epithelial-mesenchymal crosstalk that drives palatal shelf cell proliferation and outgrowth.

9. FUTURE WORK CONSIDERATIONS

My study sought to find out if *Six2* is expressed in the developing murine SP, and if it is, what role it might be playing during SP development. Although I can confidently state that all the experimental objectives of my thesis project were met, my study raises at least as many questions as it has answered. My data reveals a detailed expression analysis of *Six2* in the developing SP. Using qPCR and Western blot analyses, I demonstrate that *Six2* is expressed across all stages of SP development, while IHC reveals the spatial expression profile of this gene product in both the epithelium and mesenchyme of the developing palate. However, studies on precise *Six2* mRNA localization need to be conducted so as to consolidate the findings of this study in regard to expression analysis. This can be conducted by *in situ* hybridization studies with use of palatal tissue sections or whole mount *in situ* hybridization. I have also shown that *Six2* is mostly expressed during the earlier stages of SP development (E12.5-E13.5), suggesting that it might perform most of its functions at this earlier stage. In this study, *in vitro* cell proliferation analysis using MEPM cells showed that siRNA knockdown of *Six2* in *Hoxa2* null cells restored cell proliferation and *Cyclin D1* expression to wild type levels. This finding could be strengthened by using the palatal shelves of a *Six2* knock out mouse model instead of MEPM cell cultures. It would also be interesting to find out if the *Six2* knockout mouse has any defects in palate morphology. In addition, *in vivo* cell counting analysis using a cell proliferation marker such as Ki-67 or Brdu could be performed on coronal palate sections to directly compare cell proliferation in the *Six2* null and wild-type mice. Also, a *Six2/Hoxa2* double knock out mouse model could shed more light on the restoration to wild-type levels of cell proliferation and expression of cell cycle regulators such as *Cyclin D1* and *Cyclin D2*. A study can also be designed to

investigate if epithelial specific inactivation of *Six2* (by silencing *Six2* in palatal epithelial cell cultures) could have an effect on palate mesenchymal cell proliferation.

Furthermore, the palatal cell signalling pathways that *Six2* might be involved in are yet to be studied. The *Shh* signalling pathway is of particular interest because this gene is expressed in the oral palatal epithelium while *Six2* is also highly expressed in the oral half of the underlying palate mesenchyme. It is therefore possible that epithelial-mesenchymal interactions exist between *Shh* in the epithelium and *Six2* in the underlying mesenchyme. In addition, previous studies have revealed that the expression of *Osr1* and *Osr2*, which are some of the genes that are exclusively expressed in the oral half of the palatal mesenchyme is dependent on *Shh* signalling (Fig. 22). It would therefore be interesting to investigate if a linkage exists between *Shh*, *Six2*, *Osr1* and *Osr2*.




Figure 22: A schematic illustration of the pathways involved in O-N patterning during palatal shelf out growth.

Whereas *Pax9* is expressed across the O-N axis of the posterior palatal mesenchyme, it is expressed at higher levels in the nasal half of the palatal mesenchyme than in the oral half of the palatal mesenchyme in the anterior regions of the palate. In addition, *Pax9* has been reported to have inhibitory effects on both *Osr2* and *Fgf10* in the palate mesenchyme. *Fgf10* is exclusively expressed in the oral half of the palatal mesenchyme. *Six2* also exhibits a graded pattern of expression in the palate mesenchyme with higher expression in the oral half of the palatal mesenchyme than in the nasal half of the palate mesenchyme. *Six2* is highly expressed in the nasal palate epithelium but is devoid from the oral palatal epithelium. *Hoxa2* has been reported to have inhibitory effects on *Six2* expression. Whereas *Osr1* expression is limited to the oral half of the palate mesenchyme, *Osr2* is expressed at high levels in the oral half of the palatal mesenchyme than in the nasal half. Expression of both *Osr1* and *Osr2* is dependent on *Shh* signalling. Also, a pathway involving *Dlx5-Fgf7* signalling acts as a repressor of *Shh* signalling and controls outgrowth along the ON axis. The dotted line denotes the O-N boundary of the palatal mesenchyme. Nasal palatal epithelium (NE), Nasal half of the palatal mesenchyme (NM), Oral palatal epithelium (OE), Oral half of the palatal mesenchyme (OM), Tongue (T). Arrows indicate inductive relationships, solid lines represent direct physical interaction while blunt arrows indicate inhibitory effects. Purple boxes indicate expression in the oral half of the palate mesenchyme while orange boxes indicate expression in the nasal half of the palate mesenchyme. The size of the boxes is directly proportional to the extent of expression in the epithelium and the oral or nasal half of the palate mesenchyme.

Pax9 has been reported to have an inhibitory effect on the expression of both *Fgf10* and *Osr2*. *Pax9* mRNA has been reported to be preferentially expressed in the nasal half of the palatal mesenchyme in the anterior palate. Incidentally, my study shows that *Six2* is present only in the nasal palatal epithelium and that there are more actively proliferating cells that co-express *Six2* in the nasal half of the palate mesenchyme. Additional studies therefore can be conducted to reveal the relationship between *Six2*, *Pax9*, *Osr2* and *Fgf10* in the nasal half of the palate mesenchyme. *Fgf7* and *Dlx5* are the other genes that have been found to be expressed spatially and temporally in the developing palate. Additionally, tissue specific inactivation of *Six2* in either the epithelium or mesenchyme of the palate can be used to find out if *Six2* is acting upstream of any of these genes to regulate their expression. This study also reported that the knock down of *Six2* in MEPM cells did not have an effect on the expression of *Bmp4*, *Msx1*, *Barx1* and *Ptx1*. This could be because these genes might be acting up stream of *Six2*. Further investigations using Chromatin immunoprecipitation (ChIP) analysis can be performed to confirm whether these genes are indeed acting upstream to regulate *Six2* expression.

10. REFERENCES

- Alappat, S. R., Zhang, Z., Suzuki, K., Zhang, X., Liu, H., Jiang, R., Yamada, G. and Chen, Y. P. (2005). The cellular and molecular etiology of the cleft secondary palate in *Fgf10* mutant mice. *Dev. Biol.* 277: 102-113.
- Alwazzan M, Newman E, Hamshire MG, Brook JD, (1999). Myotonic dystrophy is associated with a reduced level of RNA from the DMWD allele adjacent to the expanded repeat. *Hum. Mol. Genet.* 8:1491-1497.
- Baek, J. A., Lan, Y., Liu, H., Maltby, K. M., Mishina, Y. and Jiang, R. (2011). *Bmpr1a* signaling plays critical roles in palatal shelf growth and palatal bone formation. *Dev. Biol.* 350, 520-531.
- Barlow, A. J., Bogardi, J. P., Ladher, R. and Francis-West, P. H. (1999). Expression of chick *Barx-1* and its differential regulation by *FGF-8* and *BMP* signaling in the maxillary primordia. *Dev. Dyn.* 214, 291-302.
- Belloni, E., Muenke, M., Roessler, E., Traverso, G., Siegel-Bartelt, J., Frumkin, A., Mitchell, H. F., Donis-Keller, H., Helms, C., Hing, A. V. (1996). Identification of Sonic hedgehog as a candidate gene responsible for holoprosencephaly. *Nat. Genet.* 14: 353-356.
- Boucher CA, King SK, Carey N, Krahe R, Winchester CL, Rahman S, Crearin T, Meghji P, Bailey MES, Chartier FL, Brown SD, Siciliano MJ, Johnson KJ, (1995). A novel homeodomain-encoding gene is associated with a large CpG island interrupted by the myotonic dystrophy unstable CTG repeat. *Hum. Mol. Genet.* 4:1919-1925.
- Boucher, C., Carey, N., Edwards, Y., Siciliano, M. and Johnson, K. (1996). Cloning of the human SIX1 gene and its assignment to chromosome 14. *Genomics* 33: 140-142

- Brodbeck, S., Besesnbeck, B. and Englert, C, (2004). The transcription factor *Six2* activates expression of the *Gdnf* gene as well as its own promoter. *Mech. Dev.* 121: 1211-1222.
- Bronner-Fraser M. (1993). Neural crest cell migration in the developing embryo. *Trends Cell Biol.* 3:392–397.
- Brown, S. A., Warburton, D., Brown, L. Y., Yu, C. Y., Roeder, E. R., Stengel-Rutkowski, S., Hennekam, R. C. and Muenke, M. (1998). Holoprosencephaly due to mutations in *ZIC2*, a homologue of *Drosophila* odd-paired. *Nat. Genetics* 20: 180-183.
- Bush, J. O. and Jiang, R. (2012). Palatogenesis: morphogenetic and molecular mechanisms of secondary palate development. *Development* 139, 231-243.
- Chai, Y. and Maxson, R.E. Jr (2006). Recent advances in craniofacial morphogenesis. *Dev. Dyn.* 235: 2353–2375.
- Cheyette BNR, Green PJ, Martin K, Garren H, Hartenstein V, Zipursky SL, (1994). The *Drosophila* sine oculis locus encodes a homeodomain-containing protein required for the development of the entire visual system. *Neuron.* 12:977-996.
- Christensen, K.L., Patrick, A.N., McCoy, E.L., Ford, H.L., (2008). The *Six* family of homeobox genes in development and cancer. *Adv. Cancer Res.* 101: 93–126.
- Christopher M Hindson, John R Chevillet, Hilary A Briggs, Emily N Gallichotte, Ingrid K Ruf, Benjamin J Hindson, Robert L Vessella & Muneesh Tewari. (2013). Absolute quantification by droplet digital PCR versus analog real-time PCR. *nmeth.* 2633,10.1038.
- Chu-An Wang, David Drasin, Catherine Pham, Paul Jedlicka, Vadym Zaberezhnyy, Michelle Guney, Howard Li, Raphael Nemenoff, James C.

Costello, Aik-Choon Tan and Heide L. Ford. (2014). Homeoprotein *Six2* promotes breast cancer metastasis via transcriptional and epigenetic control of E-Cadherin expression. *Cancer Res*, 74(24): 7357-70.

- Couly GF, Coltey PM, Le Douarin NM, (1993). The triple origin of skull in higher vertebrates — a study in quail-chick chimeras. *Development*, 117:409-429.
- Ferguson, M.W.J, (1998). Palate development. *Development*; 103(Suppl.), 41–60.
- Fitchett, J. E. and Hay, E. D. (1989). Medial edge epithelium transforms to mesenchyme after embryonic palatal shelves fuse. *Dev. Biol.* 131, 455-474.
- Fitzpatrick, D. R., Denhez, F., Kondaiah, P. and Akhurst, R. J. (1990). Differential Expression of TGF Beta Isoforms in Murine Palatogenesis. *Dev. Biol.* 109, 585-595.
- Fogelgren B., Kuroyama MC., McBratney-Owen B., Spence AA., Melahn LE., Anawati MK., Cabatbat C., Alarcon VB., Marikawa Y., and Lozanoff S, (2008). Misexpression of *Six2* is associated with heritable frontonasal dysplasia and renal hypoplasia in 3H1 *Br* mice. *Developmental Dynamics*; 237:1767-1779.
- Fogelgren B., Yang S., Sharp IC., Huckstep OJ., Ma W., Somponpun J., Carlson EC., Uyehara CFT., Lozanoff S, (2009). Deficiency in *Six2* during prenatal development is associated with reduced nephron number, chronic renal failure, and hypertension in *Br/+* adult mice. *American Journal of Physiology (Renal)*; 296(5):F1166-78.
- Gallardo ME, Lopez-Rios J, Fernaud-Espinosa I, Granadino B, Sanz R, Ramos C, Ayuso C, Seller MJ, Brunner HG, Bovolenta P, de Cordoba SR, (1999). Genomic cloning and characterization of the human homeobox

gene *Six6* reveals a cluster of *Six* genes in chromosome 14 and associates *Six6* hemizyosity with bilateral anophthalmia and pituitary anomalies. *Genomics*; 61:82-91.

- Gendron-Maguire, M., Mallo, M., Zhang, M. and Gridley, T. (1993). *Hoxa-2* mutant mice exhibit homeotic transformation of skeletal elements derived from cranial neuralcrest. *Cell*, 75:1317-1331.
- Gong, K.Q., Yallowitz, A.R., Sun, H., Dressler, G.R., Wellik, D.M., (2007). A *Hox-Eya-Pax* complex regulates early kidney developmental gene expression. *Mol. Cell. Biol*, 21:7661–7668.
- Gorlin, R.J., Cohen, M.M., Jr., and Hennekam, R.C.M. (2001). *Syndromes of the head and neck*. 4th edition. Oxford University Press. New York, New York, USA. 850–853.
- Granadino B, Gallardo ME, Lopez-Rios J *et al*, (1999): Genomic cloning, structure, expression pattern, and chromosomal location of the human *SIX3* gene. *Genomics*, 55: 100–105.
- Griffith, C.M and Hay, E.D.(1992). Epithelial-mesenchymal transformation during palatal fusion: carboxyfluorescein traces cells at light and electron microscopic levels. *Development* 116, 1087–1099.
- Gritli-Linde A, (2007). Molecular control of secondary palate development. *Dev Biol*, 301:309-326.
- Guiyuan He, Sara Tavella, Karen Piper Hanley, Michelle Self, Guillermo Oliver, Raphaëlle Grifone, Neil Hanley, Christopher Ward, Nicoletta Bobola., (2010). Inactivation of *Six2* in mouse identifies a novel genetic mechanism controlling development and growth of the cranial base. *Developmental Biology* 344: 720–730.
- Han, J., Mayo, J., Xu, X., Li, J., Bringas, P., Jr, Maas, R. L., Rubenstein, J. L. and Chai, Y. (2009). Indirect modulation of Shh signaling by *Dlx5* affects

the oral-nasal patterning of palate and rescues cleft palate in *Msx1*-null mice. *Development* 136, 4225-4233.

- Harris S, Moncrieff C, Johnson K, (1996). Myotonic dystrophy: will the real gene please step forward! *Hum Mol Genet*,5:1417±1423.
- He G, Tavella S, Hanley KP, Self M, Oliver G, Grifone R, Hanley N, Ward C, Bobola N, (2010). Inactivation of *Six2* in mouse identifies a novel genetic mechanism controlling development and growth of the cranial base. *Dev Biol*, 344:720-730.
- Heath SK, Carne S, Hoyle C Johnson, KJ, Wells DJ, (1997). Characterisation of expression of mDMAHP, a homeodomain-encoding gene at the murine DM locus. *Hum Mol Genet*, 6:651±657.
- Helms JA, Cordero D, Tapadia MD. (2005). New insights into craniofacial morphogenesis. *Development* 132:851–861.
- Hilliard, S. A., Yu, L., Gu, S., Zhang, Z. and Chen, Y. P. (2005). Regional regulation of palatal growth and patterning along the anterior-posterior axis in mice. *J. Anat.* 207, 655-667.
- Hinrichsen K. (1985). The early development of morphology and patterns of the face in the human embryo. *Adv Anat Embryol Cell Biol* 98:1–79.
- Holmes , L.B., Harvey, E.A., Coull B.A., Hunnington, K.B., Khoshbin, S., Hayes, A.M., Ryan, L.M. (2001). The teratogenicity of anticonvulsant drugs. *N Engl JMed*. 344, 1132-1138.
- Hosokawa, R., Deng, X., Takamori, K., Xu, X., Urata, M., Bringas, P., Jr and Chai, Y. (2009). Epithelial-specific requirement of *FGFR2* signaling during tooth and palate development. *J. Exp. Zool. B Mol. Dev. Evol.* 312, 343-350.

- Hu D, Helms JA. (1999). The role of sonic hedgehog in normal and abnormal craniofacial morphogenesis. *Development* 126:4873–4884.
- Hu, M. C., Mo, R., Bhella, S., Wilson, C. W., Chuang, P. T., Hui, C. C. And Rosenblum, N. D. (2006). *GLI3*-dependent transcriptional repression of *Gli1*, *Gli2* and kidney patterning genes disrupts renal morphogenesis. *Development* 133, 569-578.
- Ishibashi, M. and McMahon, A. P. (2002). A sonic hedgehog-dependent signaling relay regulates growth of diencephalic and mesencephalic primordial in the early mouse embryo. *Development* 129: 4807-4819.
- Jing Zhou, Yang Gao, Yu Lan, Shihai Jia and Rulang Jiang. (2013). *Pax9* regulates a molecular network involving *Bmp4*, *Fgf10*, *Shh* signaling and the *Osr2* transcription factor to control palate morphogenesis. *Development* 140, 4709-4718.
- Johnston MC, Bronsky PT. 1995. Prenatal craniofacial development: new insights on normal and abnormal mechanisms. *Crit Rev Oral Biol Med* 6:368–422.
- Jun-ichi Iwata, Joseph G. Hacia, Akiko Suzuki, Pedro A. Sanchez-Lara, Mark Urata, and Yang Chai. 2012. Modulation of noncanonical TGF- β signalling prevents cleft palate in *Tgfb2* mutant mice. *J Clin Invest*, 22(3):873–885.
- Kaartinen, V., Cui, X. M., Heisterkamp, N., Groffen, J. and Shuler, C. F. (1997). Transforming growth factor-beta 3 regulates transdifferentiation of medial edge epithelium during palatal fusion and associated degradation of the basement membrane. *Dev. Dyn.* 209, 255-260.
- Kaufman MH. (1992). The atlas of mouse development. New York: Academic Press. P 1–512.

- Kawakami K, Ohto H, Takizawa T, Saito T, (1996). Identification and expression of *Six* family genes in mouse retina. *FEBS Lett*, 393:259± 263.
- Kawakami, K., Sato, S., Ozaki, H. and Ikeda, K. (2000). *Six* family genes—structure and function as transcription factors and their roles in development. *BioEssays* 22: 616-626.
- Kenney, A. M. and Rowitch, D. H. (2000). Sonic hedgehog promotes G(1) cyclin expression and sustained cell cycle progression in mammalian neuronal precursors. *Mol. Cell. Biol.* 20: 9055-9067.
- Kiyoshi Kawakami, Shigeru Sato, Hidenori Ozaki, and Keiko Ikeda, (2000). *Six* family genes-structure and function as transcription factors and their roles in development. *BioEssays*; 22:616-626.
- Klesert TR, Otten AD, Bird TD, Tapscott SJ, (1997). Trinucleotide repeat expansion at the myotonic dystrophy locus reduces expression of DMAHP. *Nat Genet*, 16:402±406.
- Köntges, G. and Lumsden, A. (1996). Rhombencephalic neural crest segmentation is preserved throughout craniofacial ontogeny. *Development* 122: 3229-3242.
- Kutejova, E., Engist, B., Mallo, M., Kanzler, B. and Bobola, N, (2005). *Hoxa2* downregulates *Six2* in the neural crest derived mesenchyme. *Development*; 132: 469-478.
- Kutejova, E., Engist, B., Self, M., Oliver, G., Kirilenko, P. and Bobola, N. (2008). *Six2* functions redundantly immediately downstream of *Hoxa2*. *Development* 135: 1463-1470.
- LaBonne C, Bronner-Fraser M. (1999). Molecular mechanisms of neural crest formation. *Annu Rev Cell Dev Biol* 15:81–112.

- Lan, Y. and Jiang, R. (2009). Sonic hedgehog signaling regulates reciprocal epithelial-mesenchymal interactions controlling palatal outgrowth. *Development* 136, 1387-1396.
- Lan, Y., Ovitt, C. E., Cho, E. S., Maltby, K. M., Wang, Q. and Jiang, R. (2004). *Odd skipped related 2 (Osr2)* encodes a key intrinsic regulator of secondary palate growth and morphogenesis. *Development* 131, 3207-3216.
- Le Douarin NM, Creuzet S, Couly G, Dupin E. (2004). Neural crest cell plasticity and its limits. *Development* 131:4637–4650.
- Li, L., Lin, M., Wang, Y., Cserjesi, P., Chen, Z. and Chen, Y. (2011). *Bmpr1a* is required in mesenchymal tissue and has limited redundant function with *Bmpr1b* in tooth and palate development. *Dev. Biol.* 349, 451-461.
- Li, Q. and Ding, J. (2007). Gene expression analysis reveals that formation of the mouse anterior secondary palate involves recruitment of cells from the posterior side. *Int. J. Dev. Biol.* 51, 167-172.
- Liu, W., Lan, Y., Pauws, E., Meester-Smoor, M. A., Stanier, P., Zwarthoff, E. C. and Jiang, R. (2008). The *Mn1* transcription factor acts upstream of *Tbx22* and preferentially regulates posterior palate growth in mice. *Development* 135, 3959-3968.
- Liu, W., Sun, X., Braut, A., Mishina, Y., Behringer, R. R., Mina, M. And Martin, J. F. (2005). Distinct functions for Bmp signaling in lip and palate fusion in mice. *Development* 132, 1453-1461.
- Lobjois, V., Benazeraf, B., Bertrand, N., Medevielle, F. and Pituello, F. (2004). Specific regulation of *Cyclins D1* and *D2* by FGF and *Shh* signaling coordinates cell cycle progression, patterning, and differentiation during early steps of spinal cord development. *Dev. Biol.* 273: 195-209.

- Lozanoff S, Johnston J, Ma W, Jourdan-Le Saux C. (2001). Immunohistochemical localization of *Pax2* and associated proteins in the developing kidney of mice with renal hypoplasia. *J Histochem Cytochem* 49:1081–1097.
- Lozanoff S. (1993). Midfacial retrusion in adult Brachyrrhine mice. *Acta Anat (Basel)* 147:125–132.
- Lumsden A, Keynes R. (1989). Segmental patterns of neuronal development in the chick hindbrain. *Nature*, 337:424-428.
- Lumsden A, Sprawson N, Graham A. (1991) Segmental origin and migration of neural crest cells in the hindbrain region of the chick embryo. *Development*, 113:1281-1291.
- Lyu Z, Mao Z, Wang H, Fang Y, Chen T, Wan Q, Wang M, Wang N, Xiao J, Wei H, Li X, Liu Y and Zhou Q. (2013). *MiR-181b* targets *Six2* and inhibits the proliferation of metanephric mesenchymal cells in vitro. *Biochem Biophys Res Commun*, 440(4): 495-501.
- Ma W, Lozanoff S. (1996). Morphological deficiency in the prenatal anterior cranial base of midfacially retrognathic mice. *Journal of Anatomy*, 188:547-555.
- Ma W, Lozanoff S. (1999). Spatial and temporal distribution of cellular proliferation in the cranial base of normal and midfacially retrusive mice. *Clin Anat* 12:315–325.
- Martí'nez-A'lvarez, C., Bonelli, R., Tudela, C., Gato, A., Mena, J., O'Kane, S., Ferguson, M.W. (2000). Bulging medial edge epithelial cells and palatal fusion. *Int. J. Dev. Biol.* 44: 331– 335.
- McBratney, B.M., Margaryan, E., Ma, W., Urban, Z. and Lozanoff, S. (2003). Frontonasal dysplasia in Br/Br mice. *Anatomical Record*; 271A:291-302.

- McMahon, A. P., Ingham, P. W. and Tabin, C. J. (2003). Developmental roles and clinical significance of hedgehog signaling. *Curr. Top. Dev. Biol.* 53, 1-114.
- Meng L, Bian Z, Torensma R, Von den Hoff JW. (2009). Biological mechanisms in palatogenesis and cleft palate. *J Dent Res*, 88:22-33.
- Mill, P., Mo, R., Hu, M. C., Dagnino, L., Rosenblum, N. D. and Hui, C. C.(2005). *Shh* controls epithelial proliferation via independent pathways that converge on N-Myc. *Dev. Cell* 9: 293-303.
- Mitchell LE. (2009). Epidemiology of cleft lip and palate. In *Comprehensive Cleft Care*. Edited by: Losee JE, Kirschner RE. McGraw-Hill, Toronto, ON, Canada:35-41.
- Nawshad A. (2008). Palatal seam disintegration: to die or not to die? that is no longer the question. *Dev Dynam*, 237:2643-2656.
- Neilson KM, Pignoni F, Yan B, Moody SA. (2010). Developmental expression patterns of candidate cofactors for vertebrate six family transcription factors. *Dev Dyn*, 239: 3446-3466.
- Noden D, (1983)The role of the neural crest in patterning of avian cranial skeletal, connective, and muscle tissues. *Dev Biol*, 96:144-165.
- Ohto H, Takizawa T, Saito T, Kobayashi M, Ikeda K, Kawakami K. (1998). Tissue and developmental distribution of Six family gene products. *Int J Dev Biol*, 42:141±148.
- Oliver G, Mailhos A, Wehr R *et al*, 1995). SIX3, a murine homologue of the *sine oculis* gene, demarcates the most anterior border of the developing neural plate and is expressed during eye development. *Development*, 121: 4045–4055.

- Oliver, G., Wehr, R., Jenkins, N. A., Copeland, N. G., Cheyette, B. N., Hartenstein, V., Zipursky, S. L. and Gruss, P. (1995). Homeobox genes and connective tissue patterning. *Development*, 121: 693-705.
- Otten AD, Tapscott SJ. (1995). Triplet repeat expansion in myotonic dystrophy alters the adjacent chromatin structure. *Proc Natl Acad Sci USA*, 92:5465±5469.
- Pantalacci, S., Prochazka, J., Martin, A., Rothova, M., Lambert, A., Bernard, L., Charles, C., Viriot, L., Peterkova, R. and Laudet, V. (2008). Patterning of palatal rugae through sequential addition reveals an anterior/posterior boundary in palatal development. *BMC Dev. Biol.* 8, 116.
- Pauws, E., Moore, G. E. and Stanier, P. (2009). A functional haplotype variant in the TBX22 promoter is associated with cleft palate and ankyloglossia. *J. Med. Genet.* 46, 555-561.
- Pelton, R. W., Hogan, B. L., Miller, D. A. and Moses, H. L. (1990). Differential Expression of Genes Encoding *TGFs Beta 1, Beta 2, and Beta 3* during Murine Palate Formation. *Dev. Biol.* 141, 456-460.
- Peters, H., Neubüser, A., Kratochwil, K. and Balling, R. (1998). *Pax9*-deficient mice lack pharyngeal pouch derivatives and teeth and exhibit craniofacial and limb abnormalities. *Genes Dev.* 12, 2735-2747.
- Rice, R., Connor, E. and Rice, D. P. (2006). Expression patterns of Hedgehog signalling pathway members during mouse palate development. *Gene Expr. Patterns* 6, 206-212.
- Rice, R., Spencer-Dene, B., Connor, E. C., Gritli-Linde, A., McMahon, A. P., Dickson, C., Thesleff, I. and Rice, D. P. (2004). Disruption of *Fgf10/Fgfr2b* coordinated epithelial-mesenchymal interactions causes cleft palate. *J. Clin. Invest.* 113, 1692-1700.

- Richman JM. (1995). Head development. Craniofacial genetics makes headway. *Curr Biol*, 5:345–8.
- Rijli FM, Mark M, Lakkaraju S, Dierich A, Dolle' P, Chambon P. (1993). A homeotic transformation is generated in the rostral branchial region of the head by disruption of *Hoxa2*, which acts as a selector gene. *Cell* 75:1333–1349.
- Roessler E, Muenke M. (1998): Holoprosencephaly: a paradigm for the complex genetics of brain development. *J Inherit Metab Dis*, 21: 481–497.
- Roessler, E. and Muenke, M. (2003). How a Hedgehog might see holoprosencephaly. *Hum. Mol. Genet.* 12, R15-R25.
- Santagati F, Minoux M, Ren S-Y, Rijli FM. (2005). Temporal requirement of *Hoxa2* in cranial neural crest skeletal morphogenesis. *Development*, 132:4927–4936.
- Self, M., Lagutin, O. V., Bowling, B., Hendrix, J., Cai, Y., Dressler, G. R. And Oliver, G. (2006). *Six2* is required for suppression of nephrogenesis and progenitor renewal in the developing kidney. *EMBO J*, 25: 5214-5228.
- Selleck M.A. and Bronner-fraser M. (1995). Origins of the avian neural crest: the role of neural plate-epidermal interactions. *Development*, 121:525-538
- Seo HC, Curtiss J, Mlodzik M, Fjose A. (1999). Six class homeobox genes in drosophila belong to three distinct families and are involved in head development. *Mech Dev*, 83:127±139.
- Shuler CF. (1995). Programmed cell death and cell transformation in craniofacial development. *Crit Rev Oral Biol Med*, 6:202–17.

- Shuler, C.F., Halpern, D.E., Guo,Y., and Sank, A.C.(1992).Medial edge epithelium fate traced by cell lineage analysis during epithelial-mesenchymal transformation *in vivo*. *Dev.Biol.* 154, 318–330.
- Singh GD, Johnston J, Ma W, Lozanoff S. (1998). Cleft palate formation in fetal *Br* mice with midfacial retrusion: tenascin, fibronectin, laminin, and type IV collagen immunolocalization. *Cleft Palate Craniofac J*, 35:65–76.
- Smith TM, Wang X, Zhang W, Kulyk W, Nazarali AJ. (2009). Hoxa2 plays a direct role in murine palate development. *Dev Dynam*, 238:2364-2373.
- Stapleton, P., Weith, A., Urbánek, P., Kozmik, Z. and Busslinger, M. (1993). Chromosomal localization of seven PAX genes and cloning of a novel family member, PAX-9. *Nat. Genet.* 3, 292-298.
- Sulik KK. Dr. Beverly R. (1990). Rollnick memorial lecture. Normal and abnormal craniofacial embryogenesis. *Birth Defects Orig Artic Ser*, 26:1–18.
- Sun, D., Vanderburg, C. R., Odierana, G. S. and Hay, E. D. (1998). TGFbeta3 Promotes Transformation of Chicken Palate Medial Edge Epithelium to Mesenchyme *in Vitro*. *Development* 125, 95-105.
- Tara M.Smith, Scott Lozanoff, Paul P. Iyyanar and AdilJ.Nazarali. (2013). Molecular signalling along the anterior-posterior axis of early palate development. *Frontiers in Physiology.; Craniofacial Biology*. Volume 3, Article 488.
- Thornton CA, Wymer JP, Simmons Z, McClain C, Moxley RT. (1997). Expansion of the myotonic dystrophy CTG repeat reduces expression of the flanking DMAHP gene. *Nat Genet*, 16:407±409.
- Tyler MS, Pratt RM. (1980). Effect of epidermal growth factor on secondary palatal epithelium *in vitro*: tissue isolation and recombination studies. *J Embryol Exp Morphol*, 58: 93–106.

- Vaziri Sani F, Hallberg K, Harfe BD, Mc-Mahon AP, Linde A, Gritli-Linde A. (2005). Fate-mapping of the epithelial seam during palatal fusion rules out epithelial-mesenchymal transformation. *Dev Biol*, 285:490–495.
- Wallis DE, Roessler E, Hehr. (1999): Mutations in the homeodomain of the human *SIX3* gene cause holoprosencephaly. *Nat Genet*, 22: 196–198.
- Walsh CA. (1999). Genetic malformations of the human cerebral cortex. *Neuron*, 23:19±29.
- Wellik, D.M., Hawkes, P.J., Capecchi, M.R. (2002). Hox11 paralogous genes are essential for metanephric kidney induction. *Genes Dev*. 16, 1423–1432.
- Welsh, I. C. and O'Brien, T. P. (2009). Signaling integration in the rugae growth zone directs sequential SHH signaling center formation during the rostral outgrowth of the palate. *Dev. Biol.* 336, 53-67.
- Welsh, I. C., Hagge-Greenberg, A. and O'Brien, T. P. (2007). A Dosage-Dependent Role for *Spry2* in Growth and Patterning during Palate Development. *Mech. Develop.* 124, 746-761.
- Winchester CL, Ferrier RK, Sermoni A, Clark BJ, Johnson KJ. (1999). Characterization of the expression of DMPK and SIX5 in the human eye and implications for pathogenesis in myotonic dystrophy. *Hum Mol Genet*, 8:481±492.
- Xiaoyan Lv, Zhaomin Mao, Zhongshi Lyu, Pengzong Zhang, Alan Zhan, Jian Wang, Hui Yang, Mi Li, Honglian Wang, QianYa Wan, Hongyuan Wei, Ming Wang, Nian Wang, Xun Li, Yi Liu, Hui Zhao and Qin Zhou. (2014). MiR181c promotes apoptosis and suppresses proliferation of metanephric mesenchyme cells by targeting *Six2* in vitro. *Cell Biochem Funct*, 32: 571-579.

- Xiong, W., He, F., Morikawa, Y., Yu, X., Zhang, Z., Lan, Y., Jiang, R., Cserjesi, P. and Chen, Y. (2009). *Hand2* is required in the epithelium for palatogenesis in mice. *Dev. Biol.* 330, 131-141.
- Yu, L., Gu, S., Alappat, S., Song, Y., Yan, M., Zhang, X., Zhang, G., Jiang, Y., Zhang, Z., Zhang, Y. et al. (2005). *Shox2*-deficient mice exhibit a rare type of incomplete clefting of the secondary palate. *Development* 132, 4397-4406.
- Zhang, Y., Zhao, X., Hu, Y., St Amand, T., Zhang, M., Ramamurthy, R., Qiu, M. And Chen, Y. (1999). *Msx1* is required for the induction of Patched by Sonic hedgehog in the mammalian tooth germ. *Dev. Dyn.* 215, 45-53.
- Zhang, Z., Song, Y., Zhao, X., Zhang, X., Fermin, C. and Chen, Y. (2002). Rescue of cleft palate in *Msx1*-deficient mice by transgenic *Bmp4* reveals a network of *BMP* and *Shh* signaling in the regulation of mammalian palatogenesis. *Development* 129, 4135- 4146.

11. APPENDIX

In vivo cell counting analysis

Table 1: N of 1

Dapi positive cells (total cell number)						
	Anterior		Medial		Posterior	
	Oral	Nasal	Oral	Nasal	Oral	Nasal
Wild type	134	148	320	144	256	283
<i>Hoxa2</i> null	144	170	182	206	245	266
Ki-67 positive cells						
Wild type	49	72	108	154	157	261
<i>Hoxa2</i> null	90	120	124	173	170	228
Six2/Ki-67 double positive cells						
Wild type	22	39	56	92	130	140
<i>Hoxa2</i> null	34	43	72	105	127	91
Six2 positive cells						
Wild type	92	84	256	159	182	127
<i>Hoxa2</i> null	138	109	179	182	232	205

Table 2: N of 2

Dapi positive cells (total cell number)						
	Anterior		Medial		Posterior	
	Oral	Nasal	Oral	Nasal	Oral	Nasal
Wild type	223	262	247	250	254	291
<i>Hoxa2</i> null	236	227	202	181	247	277
Ki-67 positive cells						
Wild type	74	90	75	100	63	97
<i>Hoxa2</i> null	107	126	98	154	168	193
Six2/Ki-67 double positive cells						
Wild type	53	50	54	65	56	59
<i>Hoxa2</i> null	74	77	77	112	125	107
Six2 positive cells						
Wild type	170	145	180	154	189	171
<i>Hoxa2</i> null	200	170	196	168	241	236

Table 3: N of 3

Dapi positive cells (total cell number)						
	Anterior		Medial		Posterior	
	Oral	Nasal	Oral	Nasal	Oral	Nasal
Wild type	164	184	172	194	232	291
<i>Hoxa2</i> null	142	147	193	240	250	211
Ki-67 positive cells						
Wild type	82	95	95	101	102	118
<i>Hoxa2</i> null	83	92	82	140	120	136
Six2/Ki-67 double positive cells						
Wild type	34	42	77	78	55	70
<i>Hoxa2</i> null	35	26	65	87	82	104
Six2 positive cells						
Wild type	96	82	129	102	145	112
<i>Hoxa2</i> null	121	114	181	179	244	198

Table 4: N of 4

Dapi positive cells (total cell number)						
	Anterior		Medial		Posterior	
	Oral	Nasal	Oral	Nasal	Oral	Nasal
Wild type	220	247	193	240	203	200
<i>Hoxa2</i> null	88	123	145	160	225	208
Ki-67 positive cells						
Wild type	41	90	128	137	81	91
<i>Hoxa2</i> null	59	66	72	111	124	121
Six2/Ki-67 double positive cells						
Wild type	46	44	104	95	51	58
<i>Hoxa2</i> null	47	46	68	95	95	90
Six2 positive cells						
Wild type	165	148	144	135	155	148
<i>Hoxa2</i> null	83	94	139	132	220	178

Table 5: Ki-67 positive cells as a percentage of Dapi positive (total) cells in the whole palate.

N	Anterior		Medial		Posterior	
	Wild type	<i>Hoxa2</i> null	Wild type	<i>Hoxa2</i> null	Wild type	<i>Hoxa2</i> null
1	42.91%	66.88%	38.78%	76.55%	47.20%	93.54%
2	33.81%	50.32%	35.21%	65.80%	43.89%	68.89%
3	43.25%	60.55%	53.55%	51.27%	42.45%	70.53%
4	28.05%	59.24%	37.23%	60.00%	58.56%	72.65%

Table 6: Ki-67 positive cells as a percentage of Dapi positive (total) cells in the Wild type palate; Oral vs. Nasal.

N	Anterior		Medial		Posterior	
	Oral	Nasal	Oral	Nasal	Oral	Nasal
1	35.57%	48.65%	40.37%	52.49%	61.33%	76.87%
2	33.18%	34.35%	30.36%	40.00%	44.98%	58.99%
3	46.64%	51.63%	47.22%	52.06%	43.97%	63.45%
4	18.64%	36.44%	45.32%	54.34%	39.90%	49.19%

Table 7: Ki-67 positive cells as a percentage of Dapi positive (total) cells in the *Hoxa2* null palate; Oral vs. Nasal.

N	Anterior		Medial		Posterior	
	Oral	Nasal	Oral	Nasal	Oral	Nasal
1	43.78%	70.59%	63.13%	83.98%	69.39%	85.71%
2	45.34%	55.10%	48.51%	75.67%	68.02%	88.78%
3	58.45%	62.59%	42.49%	58.33%	48.00%	71.09%
4	47.98%	53.66%	49.66%	69.38%	55.10%	72.02%

Table 8: Six2 positive cells as a percentage of Dapi positive (total) cells in the whole palate.

N	Anterior		Medial		Posterior	
	Wild type	<i>Hoxa2</i> null	Wild type	<i>Hoxa2</i> null	Wild type	<i>Hoxa2</i> null
1	62.77%	76.23%	73.58%	81.45%	57.33%	96.28%
2	64.99%	79.91%	67.20%	89.49%	76.06%	98.87%
3	51.15%	81.31%	63.11%	87.25%	69.67%	95.88%
4	67.02%	74.89%	67.55%	84.61%	78.79%	78.99%

Table 9: Six2/Ki-67 double positive cells as a percentage of Six2 positive cells in the whole palate.

N	Anterior		Medial		Posterior	
	Wild type	<i>Hoxa2</i> null	Wild type	<i>Hoxa2</i> null	Wild type	<i>Hoxa2</i> null
1	27.79%	46.41%	35.66%	49.03%	52.51%	67.42%
2	32.70%	40.81%	35.63%	51.92%	52.32%	61.99%
3	42.70%	25.98%	48.98%	42.22%	48.64%	64.76%
4	28.75%	52.54%	42.64%	60.15%	47.35%	46.48%

Table 10: Six2/Ki-67 double positive cells as a percentage of Six2 positive cells in the Wild type palate; Oral vs Nasal.

N	Anterior		Medial		Posterior	
	Oral	Nasal	Oral	Nasal	Oral	Nasal
1	23.91%	31.24%	21.88%	57.86%	63.21%	71.41%
2	31.18%	34.48%	30.00%	42.21%	29.63%	34.50%
3	36.96%	27.86%	35.91%	44.70%	37.93%	62.50%
4	27.88%	34.13%	43.22%	42.78%	32.90%	39.19%

Table 11: Six2/Ki-67 double positive cells as a percentage of Six2 positive cells in the *Hoxa2* null palate; Oral vs Nasal.

N	Anterior		Medial		Posterior	
	Oral	Nasal	Oral	Nasal	Oral	Nasal
1	38.41%	39.45%	40.22%	57.69%	54.74%	61.75%
2	37.00%	45.29%	39.29%	61.54%	51.87%	64.55%
3	28.93%	41.42%	47.32%	48.60%	45.90%	69.95%
4	56.63%	39.79%	48.92%	65.78%	43.18%	50.56%

Absolute quantitative expression of palatal *Six2* mRNA by droplet digital PCR




Figure 1: *Six2* mRNA expression by ddPCR. Absolute quantitative expression of *Six2* mRNA during murine palatogenesis in wild type and *Hoxa2* null mice. In the wild type mice, *Six2* mRNA expression is comparably high at E12.5 and E13.5 and reduces at E14.5 before decreasing further at E15.5. In the *Hoxa2* null mice, expression is enhanced across all stages of palatogenesis. In the absence of *Hoxa2*, *Six2* mRNA is high at E12.5 and increases to a peak level at E13.5. This expression dramatically reduces at E14.5 before decreasing further at E15.5. Bars represent mean \pm SEM, $n=5$, * $p<0.05$, ** $p<0.01$.




Figure 2. Sample of the Genotyping Results of *Hoxa2* Transgenic Mice. A gel (using Red Safe) of the PCR analysis used in the genotyping of *Hoxa2* transgenic mice. The presence of a band with the wild-type primers only (A) confirms that the animal is wild-type. If a band is present only with the neomycin primers (C) this confirms the embryo is *Hoxa2* null. Presence of both bands indicates that the animal is a heterozygote (B).

Table 12: Primers used for qPCR analysis of *Msx1*, *Bmp4*, *Barx1* and *Ptx1*.

	Forward primer	Reverse primer
<i>Msx1</i>	CTCGGTGTCAAAGTGGAGG	GCTGAAGGGCAGGAGTGAA
<i>Barx1</i>	ATTGCGAGGACTGAGCGT	GACACCTGGGATTGGCTTC
<i>Bmp4</i>	AGGAAGGAGTAGATGTGAGAG	AGGGACGGAGACCAGATAC
<i>Ptx1</i>	ACCCGAGCCTTAGAAACC	GACCTGTTGGGGACAAGA
<i>Six2</i>	GCCAAGGAAAGGGAGAACAGC	GCGTCTTCTCATCCTCGGAAC

A: Six2- 314bp

B: Msx1- 153bp

C: Barx1- 175bp

D: Bmp4- 158bp

E: Ptx1- 214bp




Figure 3: An agarose gel showing the integrity of the primers that were used for PCR genotyping analysis. All bands were single with no formation of primer- dimmers.

Table of Reagent catalogue numbers

Reagent name	Company name	Calatogue number
<i>Six2</i> Taqman probes	Applied Biosystems	Mm03003557_S1
<i>Six2</i> antibody	Protein Tech	11562-A-P
<i>Msx1</i> Sybr green primers	Invitrogen	10336022
<i>Bmp4</i> Sybr green primers	Invitrogen	10336022
<i>Barx1</i> Sybr green primers	Invitrogen	10336022
<i>Ptx1</i> Sybr green primers	Invitrogen	10336022
Lipofectamine 3000	Invitrogen	1559381
<i>Six2</i> silencer select siRNA	Life Technologies	4390771
E-cadherin antibody	Sigma	U 3254
Ki-67 antibody	Affymetrix e-bioscience	13-5698-82

Letters of Permission to use images



Legal/Permissions
200 Old Tappan Road
Old Tappan, NJ 07675
Fax: 201-767-5956
Phone: 201-236-3263

July 13, 2015

PE Ref #191217

Dennis Okello
University of Saskatchewan
107 Wiggins Road
Saskatchewan
Canada
S7N5E5

Dear Dennis Okello:

You have our permission to include content from our text, ***CAMPBELL BIOLOGY, 9th Ed. by REECE, JANE B.; URRY, LISA A.; CAIN, MICHAEL L.; WASSERMAN, STEVEN A.; MINORSKY, PETER V.; JACKSON, ROBERT B.***, in your Thesis "Six2 Exhibits A Temporal-Spatial Expression Profile in the Developing Mouse Palate and Impacts Cell Proliferation During Murine Palatogenesis" for your course : Master of Science in Pharmacy at UNIVERSITY OF SASKATCHEWAN.

Content to be included is:

P. 1032 Figure 47.13 Neurulation in a frog embryo(a) Neural plate formation (b) Neural tube formation (c) Somites

***With Exception to Photograph in 47.13(a) upper left corner figure- listed in the photo credits page as Huw Williams and Photograph 47.13 (c) upper right corner-listed in the photo credits as Thomas Poole, Suny Health Science Center. For permission to use these two photographs, contact the source given.**

Permission is granted for use of the requested figure (excluding the photographs) to be printed for yourself, the Instructor, School and School Committee. The University of Saskatchewan also has permission to electronically store your Thesis

Please credit our material as follows:

REECE, JANE B.; URRY, LISA A.; CAIN, MICHAEL L.; WASSERMAN, STEVEN A.; MINORSKY, PETER V.; JACKSON, ROBERT B., CAMPBELL BIOLOGY, 9th Edition, © 2011. Reprinted and Electronically reproduced by permission of Pearson Education, Inc., Upper Saddle River, NJ

Sincerely,
Mary Ann Vass, Permissions Specialist

JOHN WILEY AND SONS LICENSE TERMS AND CONDITIONS

Aug 25, 2015

This Agreement between Dennis Okello ("You") and John Wiley and Sons ("John Wiley and Sons") consists of your license details and the terms and conditions provided by John Wiley and Sons and Copyright Clearance Center.

License Number	3666601139614
License date	Jul 12, 2015
Licensed Content Publisher	John Wiley and Sons
Licensed Content Publication	BioEssays
Licensed Content Title	Six family genes—structure and function as transcription factors and their roles in development
Licensed Content Author	Kiyoshi Kawakami, Shigeru Sato, Hidenori Ozaki, Keiko Ikeda
Licensed Content Date	Jun 23, 2000
Pages	11
Type of use	Dissertation/Thesis
Requestor type	University/Academic
Format	Print and electronic
Portion	Figure/table
Number of figures/tables	1
Original Wiley figure/table number(s)	Figure 3: Expression patterns and chromosomal locations of mouse Six family genes.
Will you be translating?	No
Title of your thesis / dissertation	Six2 EXHIBITS A TEMPORAL-SPATIAL EXPRESSION PROFILE IN THE DEVELOPING MOUSE PALATE AND IMPACTS CELL PROLIFERATION DURING MURINE PALATOGENESIS
Expected completion date	Jul 2015
Expected size (number of pages)	240
	Dennis Okello 107 Wiggins Road
Requestor Location	Saskatoon, SK, SK S7N 5E5 Canada Attn: Dennis Okello
Billing Type	Invoice Dennis Okello 107 Wiggins Road
Billing Address	Saskatoon, SK, SK S7N 5E5 Canada


Total

Terms and Conditions

Attn: Dennis Okello

0.00 CAD

TERMS AND CONDITIONS

This copyrighted material is owned by or exclusively licensed to John Wiley & Sons, Inc. or one of its group companies (each a "Wiley Company") or handled on behalf of a society with which a Wiley Company has exclusive publishing rights in relation to a particular work (collectively "WILEY"). By clicking  in connection with completing this licensing transaction, you agree that the following terms and conditions apply to this transaction (along with the billing and payment terms and conditions established by the Copyright Clearance Center Inc., ("CCC's Billing and Payment terms and conditions"), at the time that you opened your Rightslink account (these are available at any time at <http://myaccount.copyright.com/>).

Terms and Conditions

- The materials you have requested permission to reproduce or reuse (the "Wiley Materials") are protected by copyright.
- You are hereby granted a personal, non-exclusive, non-sub licensable (on a stand-alone basis), non-transferable, worldwide, limited license to reproduce the Wiley Materials for the purpose specified in the licensing process. This license is for a one-time use only and limited to any maximum distribution number specified in the license. The first instance of republication or reuse granted by this licence must be completed within two years of the date of the grant of this licence (although copies prepared before the end date may be distributed thereafter). The Wiley Materials shall not be used in any other manner or for any other purpose, beyond what is granted in the license. Permission is granted subject to an appropriate acknowledgement given to the author, title of the material/book/journal and the publisher. You shall also duplicate the copyright notice that appears in the Wiley publication in your use of the Wiley Material. Permission is also granted on the understanding that nowhere in the text is a previously published source acknowledged for all or part of this Wiley Material. Any third party content is expressly excluded from this permission.
- With respect to the Wiley Materials, all rights are reserved. Except as expressly granted by the terms of the license, no part of the Wiley Materials may be copied, modified, adapted (except for minor reformatting required by the new Publication), translated, reproduced, transferred or distributed, in any form or by any means, and no derivative works may be made based on the Wiley Materials without the prior permission of the respective copyright owner. You may not alter, remove or suppress in any manner any copyright, trademark or other notices displayed by the Wiley Materials. You may not license, rent, sell, loan, lease, pledge, offer as security, transfer or assign the Wiley Materials on a stand-alone basis, or any of the rights granted to you hereunder to any other person.
- The Wiley Materials and all of the intellectual property rights therein shall at all times remain the exclusive property of John Wiley & Sons Inc, the Wiley Companies, or their respective licensors, and your interest therein is only that of having possession of and the right to reproduce the Wiley Materials pursuant to Section 2 herein during the

continuance of this Agreement. You agree that you own no right, title or interest in or to the Wiley Materials or any of the intellectual property rights therein. You shall have no rights hereunder other than the license as provided for above in Section 2. No right, license or interest to any trademark, trade name, service mark or other branding ("Marks") of WILEY or its licensors is granted hereunder, and you agree that you shall not assert any such right, license or interest with respect thereto.

- NEITHER WILEY NOR ITS LICENSORS MAKES ANY WARRANTY OR REPRESENTATION OF ANY KIND TO YOU OR ANY THIRD PARTY, EXPRESS, IMPLIED OR STATUTORY, WITH RESPECT TO THE MATERIALS OR THE ACCURACY OF ANY INFORMATION CONTAINED IN THE MATERIALS, INCLUDING, WITHOUT LIMITATION, ANY IMPLIED WARRANTY OF MERCHANTABILITY, ACCURACY, SATISFACTORY QUALITY, FITNESS FOR A PARTICULAR PURPOSE, USABILITY, INTEGRATION OR NON-INFRINGEMENT AND ALL SUCH WARRANTIES ARE HEREBY EXCLUDED BY WILEY AND ITS LICENSORS AND WAIVED BY YOU
- WILEY shall have the right to terminate this Agreement immediately upon breach of this Agreement by you.
- You shall indemnify, defend and hold harmless WILEY, its Licensors and their respective directors, officers, agents and employees, from and against any actual or threatened claims, demands, causes of action or proceedings arising from any breach of this Agreement by you.
- IN NO EVENT SHALL WILEY OR ITS LICENSORS BE LIABLE TO YOU OR ANY OTHER PARTY OR ANY OTHER PERSON OR ENTITY FOR ANY SPECIAL, CONSEQUENTIAL, INCIDENTAL, INDIRECT, EXEMPLARY OR PUNITIVE DAMAGES, HOWEVER CAUSED, ARISING OUT OF OR IN CONNECTION WITH THE DOWNLOADING, PROVISIONING, VIEWING OR USE OF THE MATERIALS REGARDLESS OF THE FORM OF ACTION, WHETHER FOR BREACH OF CONTRACT, BREACH OF WARRANTY, TORT, NEGLIGENCE, INFRINGEMENT OR OTHERWISE (INCLUDING, WITHOUT LIMITATION, DAMAGES BASED ON LOSS OF PROFITS, DATA, FILES, USE, BUSINESS OPPORTUNITY OR CLAIMS OF THIRD PARTIES), AND WHETHER OR NOT THE PARTY HAS BEEN ADVISED OF THE POSSIBILITY OF SUCH DAMAGES. THIS LIMITATION SHALL APPLY NOTWITHSTANDING ANY FAILURE OF ESSENTIAL PURPOSE OF ANY LIMITED REMEDY PROVIDED HEREIN.
- Should any provision of this Agreement be held by a court of competent jurisdiction to be illegal, invalid, or unenforceable, that provision shall be deemed amended to achieve as nearly as possible the same economic effect as the original provision, and the legality, validity and enforceability of the remaining provisions of this Agreement shall not be affected or impaired thereby.
- The failure of either party to enforce any term or condition of this Agreement shall not constitute a waiver of either party's right to enforce each and every term and condition of this Agreement. No breach under this agreement shall be deemed waived or excused by either party unless such waiver or consent is in writing signed by the party granting such waiver or consent. The waiver by or consent of a party to a breach of any provision of this Agreement shall not operate or be construed as a waiver of or consent to any other or subsequent breach by such other party.
- This Agreement may not be assigned (including by operation of law or otherwise) by you without WILEY's prior written consent.
- Any fee required for this permission shall be non-refundable after thirty (30) days from

receipt by the CCC.

- These terms and conditions together with CCC's Billing and Payment terms and conditions (which are incorporated herein) form the entire agreement between you and WILEY concerning this licensing transaction and (in the absence of fraud) supersedes all prior agreements and representations of the parties, oral or written. This Agreement may not be amended except in writing signed by both parties. This Agreement shall be binding upon and inure to the benefit of the parties' successors, legal representatives, and authorized assigns.
- In the event of any conflict between your obligations established by these terms and conditions and those established by CCC's Billing and Payment terms and conditions, these terms and conditions shall prevail.
- WILEY expressly reserves all rights not specifically granted in the combination of (i) the license details provided by you and accepted in the course of this licensing transaction, (ii) these terms and conditions and (iii) CCC's Billing and Payment terms and conditions.
- This Agreement will be void if the Type of Use, Format, Circulation, or Requestor Type was misrepresented during the licensing process.
- This Agreement shall be governed by and construed in accordance with the laws of the State of New York, USA, without regards to such state's conflict of law rules. Any legal action, suit or proceeding arising out of or relating to these Terms and Conditions or the breach thereof shall be instituted in a court of competent jurisdiction in New York County in the State of New York in the United States of America and each party hereby consents and submits to the personal jurisdiction of such court, waives any objection to venue in such court and consents to service of process by registered or certified mail, return receipt requested, at the last known address of such party.

WILEY OPEN ACCESS TERMS AND CONDITIONS

Wiley Publishes Open Access Articles in fully Open Access Journals and in Subscription journals offering Online Open. Although most of the fully Open Access journals publish open access articles under the terms of the Creative Commons Attribution (CC BY) License only, the subscription journals and a few of the Open Access Journals offer a choice of Creative Commons Licenses:: Creative Commons Attribution (CC-BY) license [Creative Commons Attribution Non-Commercial \(CC-BY-NC\) license](#) and [Creative Commons Attribution Non-Commercial-NoDerivs \(CC-BY-NC-ND\) License](#). The license type is clearly identified on the article.

Copyright in any research article in a journal published as Open Access under a Creative Commons License is retained by the author(s). Authors grant Wiley a license to publish the article and identify itself as the original publisher. Authors also grant any third party the right to use the article freely as long as its integrity is maintained and its original authors, citation details and publisher are identified as follows: [Title of Article/Author/Journal Title and Volume/Issue. Copyright (c) [year] [copyright owner as specified in the Journal]. Links to the final article on Wiley's website are encouraged where applicable.

The Creative Commons Attribution License

The [Creative Commons Attribution License \(CC-BY\)](#) allows users to copy, distribute and transmit an article, adapt the article and make commercial use of the article. The CC-BY license permits commercial and non-commercial re-use of an open access article, as long as the author is properly attributed.

The Creative Commons Attribution License does not affect the moral rights of authors, including without limitation the right not to have their work subjected to derogatory treatment. It also does not affect any other rights held by authors or third parties in the article, including without limitation the rights of privacy and publicity. Use of the article must not assert or imply, whether implicitly or explicitly, any connection with, endorsement or sponsorship of such use by the author, publisher or any other party associated with the article.

For any reuse or distribution, users must include the copyright notice and make clear to others that the article is made available under a Creative Commons Attribution license, linking to the relevant Creative Commons web page.

To the fullest extent permitted by applicable law, the article is made available as is and without representation or warranties of any kind whether express, implied, statutory or otherwise and including, without limitation, warranties of title, merchantability, fitness for a particular purpose, non-infringement, absence of defects, accuracy, or the presence or absence of errors.

Creative Commons Attribution Non-Commercial License

The [Creative Commons Attribution Non-Commercial \(CC-BY-NC\) License](#) permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.(see below)

Creative Commons Attribution-Non-Commercial-NoDerivs License

The [Creative Commons Attribution Non-Commercial-NoDerivs License](#) (CC-BY-NC-ND) permits use, distribution and reproduction in any medium, provided the original work is properly cited, is not used for commercial purposes and no modifications or adaptations are made. (see below)

Use by non-commercial users

For non-commercial and non-promotional purposes, individual users may access, download, copy, display and redistribute to colleagues Wiley Open Access articles, as well as adapt, translate, text- and data-mine the content subject to the following conditions:

- The authors' moral rights are not compromised. These rights include the right of "paternity" (also known as "attribution" - the right for the author to be identified as such) and "integrity" (the right for the author not to have the work altered in such a way that the author's reputation or integrity may be impugned).
- Where content in the article is identified as belonging to a third party, it is the obligation of the user to ensure that any reuse complies with the copyright policies of the owner of that

content.

- If article content is copied, downloaded or otherwise reused for non-commercial research and education purposes, a link to the appropriate bibliographic citation (authors, journal, article title, volume, issue, page numbers, DOI and the link to the definitive published version on **Wiley Online Library**) should be maintained. Copyright notices and disclaimers must not be deleted.
- Any translations, for which a prior translation agreement with Wiley has not been agreed, must prominently display the statement: "This is an unofficial translation of an article that appeared in a Wiley publication. The publisher has not endorsed this translation."

Use by commercial "for-profit" organisations

Use of Wiley Open Access articles for commercial, promotional, or marketing purposes requires further explicit permission from Wiley and will be subject to a fee. Commercial purposes include:

- Copying or downloading of articles, or linking to such articles for further redistribution, sale or licensing;
- Copying, downloading or posting by a site or service that incorporates advertising with such content;
- The inclusion or incorporation of article content in other works or services (other than normal quotations with an appropriate citation) that is then available for sale or licensing, for a fee (for example, a compilation produced for marketing purposes, inclusion in a sales pack)
- Use of article content (other than normal quotations with appropriate citation) by for-profit organisations for promotional purposes
- Linking to article content in e-mails redistributed for promotional, marketing or educational purposes;
- Use for the purposes of monetary reward by means of sale, resale, licence, loan, transfer or other form of commercial exploitation such as marketing products
- Print reprints of Wiley Open Access articles can be purchased from:
<mailto:corporatesales@wiley.com>

Further details can be found on Wiley Online Library
<http://olabout.wiley.com/WileyCDA/Section/id-410895.html>

Other Terms and Conditions:

v1.9

Questions? customer care@copyright.com or +1-855-239-3415 (toll free in the US) or +1-978-646-2777.

ELSEVIER LICENSE TERMS AND CONDITIONS

Aug 25, 2015

This is a License Agreement between Dennis Okello ("You") and Elsevier ("Elsevier") provided by Copyright Clearance Center ("CCC"). The license consists of your order details, the terms and conditions provided by Elsevier, and the payment terms and conditions.

All payments must be made in full to CCC. For payment instructions, please see information listed at the bottom of this form.

Supplier	Elsevier Limited The Boulevard, Langford Lane Kidlington, Oxford, OX5 1GB, UK
Registered Company Number	1982084
Customer name	Dennis Okello
Customer address	107 Wiggins Road Saskatoon, SK, SK S7N 5E5
License number	3665720037925
License date	Jul 11, 2015
Licensed content publisher	Elsevier
Licensed content publication	Genomics
Licensed content title	Cloning of the HumanSIX1Gene and Its Assignment to Chromosome 14
Licensed content author	Catherine A. Boucher, Nessa Carey, Yvonne H. Edwards, Michael J. Siciliano, Keith J. Johnson
Licensed content date	1 April 1996
Licensed content volume number	33
Licensed content issue number	1
Number of pages	3
Start Page	140
End Page	142
Type of Use	reuse in a thesis/dissertation
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
Format	both print and electronic
Are you the author of this Elsevier article?	No
Will you be translating?	No
Original figure numbers	Figure 1B
Title of your thesis/dissertation	Six2 EXHIBITS A TEMPORAL-SPATIAL EXPRESSION PROFILE IN THE DEVELOPING MOUSE PALATE AND IMPACTS CELL PROLIFERATION DURING MURINE PALATOGENESIS

Expected completion date	Jul 2015
Estimated size (number of pages)	240
Elsevier VAT number	GB 494 6272 12
Permissions price	0.00 CAD
VAT/Local Sales Tax	0.00 CAD / 0.00 GBP
Total	0.00 CAD
Terms and Conditions	

INTRODUCTION

1. The publisher for this copyrighted material is Elsevier. By clicking "accept" in connection with completing this licensing transaction, you agree that the following terms and conditions apply to this transaction (along with the Billing and Payment terms and conditions established by Copyright Clearance Center, Inc. ("CCC"), at the time that you opened your Rightslink account and that are available at any time at <http://myaccount.copyright.com/>).

GENERAL TERMS

2. Elsevier hereby grants you permission to reproduce the aforementioned material subject to the terms and conditions indicated.

3. Acknowledgement: If any part of the material to be used (for example, figures) has appeared in our publication with credit or acknowledgement to another source, permission must also be sought from that source. If such permission is not obtained then that material may not be included in your publication/copies. Suitable acknowledgement to the source must be made, either as a footnote or in a reference list at the end of your publication, as follows:

"Reprinted from Publication title, Vol /edition number, Author(s), Title of article / title of chapter, Pages No., Copyright (Year), with permission from Elsevier [OR APPLICABLE SOCIETY COPYRIGHT OWNER]." Also Lancet special credit - "Reprinted from The Lancet, Vol. number, Author(s), Title of article, Pages No., Copyright (Year), with permission from Elsevier."

4. Reproduction of this material is confined to the purpose and/or media for which permission is hereby given.

5. Altering/Modifying Material: Not Permitted. However figures and illustrations may be altered/adapted minimally to serve your work. Any other abbreviations, additions, deletions and/or any other alterations shall be made only with prior written authorization of Elsevier Ltd. (Please contact Elsevier at permissions@elsevier.com)

6. If the permission fee for the requested use of our material is waived in this instance, please be advised that your future requests for Elsevier materials may attract a fee.

7. Reservation of Rights: Publisher reserves all rights not specifically granted in the combination of (i) the license details provided by you and accepted in the course of this licensing transaction, (ii) these terms and conditions and (iii) CCC's Billing and Payment

terms and conditions.

8. License Contingent Upon Payment: While you may exercise the rights licensed immediately upon issuance of the license at the end of the licensing process for the transaction, provided that you have disclosed complete and accurate details of your proposed use, no license is finally effective unless and until full payment is received from you (either by publisher or by CCC) as provided in CCC's Billing and Payment terms and conditions. If full payment is not received on a timely basis, then any license preliminarily granted shall be deemed automatically revoked and shall be void as if never granted. Further, in the event that you breach any of these terms and conditions or any of CCC's Billing and Payment terms and conditions, the license is automatically revoked and shall be void as if never granted. Use of materials as described in a revoked license, as well as any use of the materials beyond the scope of an unrevoked license, may constitute copyright infringement and publisher reserves the right to take any and all action to protect its copyright in the materials.

9. Warranties: Publisher makes no representations or warranties with respect to the licensed material.

10. Indemnity: You hereby indemnify and agree to hold harmless publisher and CCC, and their respective officers, directors, employees and agents, from and against any and all claims arising out of your use of the licensed material other than as specifically authorized pursuant to this license.

11. No Transfer of License: This license is personal to you and may not be sublicensed, assigned, or transferred by you to any other person without publisher's written permission.

12. No Amendment Except in Writing: This license may not be amended except in a writing signed by both parties (or, in the case of publisher, by CCC on publisher's behalf).

13. Objection to Contrary Terms: Publisher hereby objects to any terms contained in any purchase order, acknowledgment, check endorsement or other writing prepared by you, which terms are inconsistent with these terms and conditions or CCC's Billing and Payment terms and conditions. These terms and conditions, together with CCC's Billing and Payment terms and conditions (which are incorporated herein), comprise the entire agreement between you and publisher (and CCC) concerning this licensing transaction. In the event of any conflict between your obligations established by these terms and conditions and those established by CCC's Billing and Payment terms and conditions, these terms and conditions shall control.

14. Revocation: Elsevier or Copyright Clearance Center may deny the permissions described in this License at their sole discretion, for any reason or no reason, with a full refund payable to you. Notice of such denial will be made using the contact information provided by you. Failure to receive such notice will not alter or invalidate the denial. In no event will Elsevier or Copyright Clearance Center be responsible or liable for any costs, expenses or damage incurred by you as a result of a denial of your permission request, other than a refund of the amount(s) paid by you to Elsevier and/or Copyright Clearance Center for denied permissions.

LIMITED LICENSE

The following terms and conditions apply only to specific license types:

15. Translation: This permission is granted for non-exclusive world **English** rights only unless your license was granted for translation rights. If you licensed translation rights you may only translate this content into the languages you requested. A professional translator must perform all translations and reproduce the content word for word preserving the integrity of the article. If this license is to re-use 1 or 2 figures then permission is granted for non-exclusive world rights in all languages.

16. Posting licensed content on any Website: The following terms and conditions apply as follows: Licensing material from an Elsevier journal: All content posted to the web site must maintain the copyright information line on the bottom of each image; A hyper-text must be included to the Homepage of the journal from which you are licensing at <http://www.sciencedirect.com/science/journal/xxxxx> or the Elsevier homepage for books at <http://www.elsevier.com/>; Central Storage: This license does not include permission for a scanned version of the material to be stored in a central repository such as that provided by Heron/XanEdu.

Licensing material from an Elsevier book: A hyper-text link must be included to the Elsevier homepage at <http://www.elsevier.com/>. All content posted to the web site must maintain the copyright information line on the bottom of each image.

Posting licensed content on Electronic reserve: In addition to the above the following clauses are applicable: The web site must be password-protected and made available only to bona fide students registered on a relevant course. This permission is granted for 1 year only. You may obtain a new license for future website posting.

17. For journal authors: the following clauses are applicable in addition to the above:

Preprints:

A preprint is an author's own write-up of research results and analysis, it has not been peer-reviewed, nor has it had any other value added to it by a publisher (such as formatting, copyright, technical enhancement etc.).

Authors can share their preprints anywhere at any time. Preprints should not be added to or enhanced in any way in order to appear more like, or to substitute for, the final versions of articles however authors can update their preprints on arXiv or RePEc with their Accepted Author Manuscript (see below).

If accepted for publication, we encourage authors to link from the preprint to their formal publication via its DOI. Millions of researchers have access to the formal publications on ScienceDirect, and so links will help users to find, access, cite and use the best available version. Please note that Cell Press, The Lancet and some society-owned have different preprint policies. Information on these policies is available on the journal homepage.

Accepted Author Manuscripts: An accepted author manuscript is the manuscript of an

article that has been accepted for publication and which typically includes author-incorporated changes suggested during submission, peer review and editor-author communications.

Authors can share their accepted author manuscript:

- <!--[if !supportLists]-->– <!--[endif]-->immediately
 - <!--[if !supportLists]--><!--[endif]-->via their non-commercial person homepage or blog
 - <!--[if !supportLists]--><!--[endif]-->by updating a preprint in arXiv or RePEc with the accepted manuscript
 - <!--[if !supportLists]--><!--[endif]-->via their research institute or institutional repository for internal institutional uses or as part of an invitation-only research collaboration work-group
 - <!--[if !supportLists]--><!--[endif]-->directly by providing copies to their students or to research collaborators for their personal use
 - <!--[if !supportLists]--><!--[endif]-->for private scholarly sharing as part of an invitation-only work group on commercial sites with which Elsevier has an agreement
- <!--[if !supportLists]-->– <!--[endif]-->after the embargo period
 - <!--[if !supportLists]--><!--[endif]-->via non-commercial hosting platforms such as their institutional repository
 - <!--[if !supportLists]--><!--[endif]-->via commercial sites with which Elsevier has an agreement

In all cases accepted manuscripts should:

- <!--[if !supportLists]-->– <!--[endif]-->link to the formal publication via its DOI
- <!--[if !supportLists]-->– <!--[endif]-->bear a CC-BY-NC-ND license - this is easy to do
- <!--[if !supportLists]-->– <!--[endif]-->if aggregated with other manuscripts, for example in a repository or other site, be shared in alignment with our hosting policy not be added to or enhanced in any way to appear more like, or to substitute for, the published journal article.

Published journal article (JPA): A published journal article (PJA) is the definitive final record of published research that appears or will appear in the journal and embodies all value-adding publishing activities including peer review co-ordination, copy-editing, formatting, (if relevant) pagination and online enrichment.

Policies for sharing publishing journal articles differ for subscription and gold open access articles:

Subscription Articles: If you are an author, please share a link to your article rather than the full-text. Millions of researchers have access to the formal publications on ScienceDirect, and so links will help your users to find, access, cite, and use the best available version.

Theses and dissertations which contain embedded PJAs as part of the formal submission

can be posted publicly by the awarding institution with DOI links back to the formal publications on ScienceDirect.

If you are affiliated with a library that subscribes to ScienceDirect you have additional private sharing rights for others' research accessed under that agreement. This includes use for classroom teaching and internal training at the institution (including use in course packs and courseware programs), and inclusion of the article for grant funding purposes.

Gold Open Access Articles: May be shared according to the author-selected end-user license and should contain a [CrossMark logo](#), the end user license, and a DOI link to the formal publication on ScienceDirect.

Please refer to Elsevier's [posting policy](#) for further information.

18. For book authors the following clauses are applicable in addition to the above: Authors are permitted to place a brief summary of their work online only. You are not allowed to download and post the published electronic version of your chapter, nor may you scan the printed edition to create an electronic version. **Posting to a repository:** Authors are permitted to post a summary of their chapter only in their institution's repository.

19. Thesis/Dissertation: If your license is for use in a thesis/dissertation your thesis may be submitted to your institution in either print or electronic form. Should your thesis be published commercially, please reapply for permission. These requirements include permission for the Library and Archives of Canada to supply single copies, on demand, of the complete thesis and include permission for Proquest/UMI to supply single copies, on demand, of the complete thesis. Should your thesis be published commercially, please reapply for permission. Theses and dissertations which contain embedded PJAs as part of the formal submission can be posted publicly by the awarding institution with DOI links back to the formal publications on ScienceDirect.

Elsevier Open Access Terms and Conditions

You can publish open access with Elsevier in hundreds of open access journals or in nearly 2000 established subscription journals that support open access publishing. Permitted third party re-use of these open access articles is defined by the author's choice of Creative Commons user license. See our [open access license policy](#) for more information.

Terms & Conditions applicable to all Open Access articles published with Elsevier:

Any reuse of the article must not represent the author as endorsing the adaptation of the article nor should the article be modified in such a way as to damage the author's honour or reputation. If any changes have been made, such changes must be clearly indicated.

The author(s) must be appropriately credited and we ask that you include the end user license and a DOI link to the formal publication on ScienceDirect.

If any part of the material to be used (for example, figures) has appeared in our publication

with credit or acknowledgement to another source it is the responsibility of the user to ensure their reuse complies with the terms and conditions determined by the rights holder.

Additional Terms & Conditions applicable to each Creative Commons user license:

CC BY: The CC-BY license allows users to copy, to create extracts, abstracts and new works from the Article, to alter and revise the Article and to make commercial use of the Article (including reuse and/or resale of the Article by commercial entities), provided the user gives appropriate credit (with a link to the formal publication through the relevant DOI), provides a link to the license, indicates if changes were made and the licensor is not represented as endorsing the use made of the work. The full details of the license are available at <http://creativecommons.org/licenses/by/4.0>.

CC BY NC SA: The CC BY-NC-SA license allows users to copy, to create extracts, abstracts and new works from the Article, to alter and revise the Article, provided this is not done for commercial purposes, and that the user gives appropriate credit (with a link to the formal publication through the relevant DOI), provides a link to the license, indicates if changes were made and the licensor is not represented as endorsing the use made of the work. Further, any new works must be made available on the same conditions. The full details of the license are available at <http://creativecommons.org/licenses/by-nc-sa/4.0>.

CC BY NC ND: The CC BY-NC-ND license allows users to copy and distribute the Article, provided this is not done for commercial purposes and further does not permit distribution of the Article if it is changed or edited in any way, and provided the user gives appropriate credit (with a link to the formal publication through the relevant DOI), provides a link to the license, and that the licensor is not represented as endorsing the use made of the work. The full details of the license are available at <http://creativecommons.org/licenses/by-nc-nd/4.0>. Any commercial reuse of Open Access articles published with a CC BY NC SA or CC BY NC ND license requires permission from Elsevier and will be subject to a fee.

Commercial reuse includes:

- <!--[if !supportLists]--> <!--[endif]--> Associating advertising with the full text of the Article
- <!--[if !supportLists]--> <!--[endif]--> Charging fees for document delivery or access
- <!--[if !supportLists]--> <!--[endif]--> Article aggregation
- <!--[if !supportLists]--> <!--[endif]--> Systematic distribution via e-mail lists or share buttons

Posting or linking by commercial companies for use by customers of those companies.

20. Other Conditions:

v1.7

Questions? customercare@copyright.com or +1-855-239-3415 (toll free in the US) or +1-978-646-2777.

NATURE PUBLISHING GROUP LICENSE TERMS AND CONDITIONS

Aug 25, 2015

This is a License Agreement between Dennis Okello ("You") and Nature Publishing Group ("Nature Publishing Group") provided by Copyright Clearance Center ("CCC"). The license consists of your order details, the terms and conditions provided by Nature Publishing Group, and the payment terms and conditions.

All payments must be made in full to CCC. For payment instructions, please see information listed at the bottom of this form.

License Number	3663431408355
License date	Jul 07, 2015
Licensed content publisher	Nature Publishing Group
Licensed content publication	Nature Reviews Neuroscience
Licensed content title	Neural crest specification: migrating into genomics
Licensed content author	Laura S. Gammill and Marianne Bronner-Fraser
Licensed content date	Oct 1, 2003
Volume number	4
Issue number	10
Type of Use	reuse in a dissertation / thesis
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
High-res required	no
Figures	Figure 1: Border induction and Neurulation
Author of this NPG article	no
Your reference number	None
Title of your thesis / dissertation	Six2 EXHIBITS A TEMPORAL-SPATIAL EXPRESSION PROFILE IN THE DEVELOPING MOUSE PALATE AND IMPACTS CELL PROLIFERATION DURING MURINE PALATOGENESIS
Expected completion date	Jul 2015
Estimated size (number of pages)	140
Total	0.00 CAD
Terms and Conditions	

Terms and Conditions for Permissions

Nature Publishing Group hereby grants you a non-exclusive license to reproduce this

material for this purpose, and for no other use, subject to the conditions below:

1. NPG warrants that it has, to the best of its knowledge, the rights to license reuse of this material. However, you should ensure that the material you are requesting is original to Nature Publishing Group and does not carry the copyright of another entity (as credited in the published version). If the credit line on any part of the material you have requested indicates that it was reprinted or adapted by NPG with permission from another source, then you should also seek permission from that source to reuse the material.
2. Permission granted free of charge for material in print is also usually granted for any electronic version of that work, provided that the material is incidental to the work as a whole and that the electronic version is essentially equivalent to, or substitutes for, the print version. Where print permission has been granted for a fee, separate permission must be obtained for any additional, electronic reuse (unless, as in the case of a full paper, this has already been accounted for during your initial request in the calculation of a print run). NB: In all cases, web-based use of full-text articles must be authorized separately through the 'Use on a Web Site' option when requesting permission.
3. Permission granted for a first edition does not apply to second and subsequent editions and for editions in other languages (except for signatories to the STM Permissions Guidelines, or where the first edition permission was granted for free).
4. Nature Publishing Group's permission must be acknowledged next to the figure, table or abstract in print. In electronic form, this acknowledgement must be visible at the same time as the figure/table/abstract, and must be hyperlinked to the journal's homepage.
5. The credit line should read:
Reprinted by permission from Macmillan Publishers Ltd: [JOURNAL NAME] (reference citation), copyright (year of publication)
For AOP papers, the credit line should read:
Reprinted by permission from Macmillan Publishers Ltd: [JOURNAL NAME], advance online publication, day month year (doi: 10.1038/sj.[JOURNAL ACRONYM].XXXXX)

Note: For republication from the *British Journal of Cancer*, the following credit lines apply.

Reprinted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication)
For AOP papers, the credit line should read:
Reprinted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME], advance online publication, day month year (doi: 10.1038/sj.[JOURNAL ACRONYM].XXXXX)

6. Adaptations of single figures do not require NPG approval. However, the adaptation should be credited as follows:

Adapted by permission from Macmillan Publishers Ltd: [JOURNAL NAME] (reference citation), copyright (year of publication)

Note: For adaptation from the *British Journal of Cancer*, the following credit line applies.

Adapted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication)

7. Translations of 401 words up to a whole article require NPG approval. Please visit <http://www.macmillanmedicalcommunications.com/> for more information. Translations of up to a 400 words do not require NPG approval. The translation should be credited as follows:

Translated by permission from Macmillan Publishers Ltd: [JOURNAL NAME] (reference citation), copyright (year of publication).

Note: For translation from the *British Journal of Cancer*, the following credit line applies.

Translated by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication)

We are certain that all parties will benefit from this agreement and wish you the best in the use of this material. Thank you.

Special Terms:

v1.1

Questions? customercare@copyright.com or +1-855-239-3415 (toll free in the US) or +1-978-646-2777.



Okello, Dennis Okori

Fri 7/10/2015 8:58 PM

Sent Items

← REPLY

↩ REPLY ALL

→ FORWARD



Mark as unread

To: permissions <permissions@biologists.com>;

Hello Sir/Madam,

I am a Master of Science student in the college of Pharmacy and Nutrition at the University of Saskatchewan, Canada. I am writing to request for permission to use Figure 1 from the article: **Helms JA, Cordero D, Tapadia MD. (2005). New insights into craniofacial morphogenesis. Development 132:851–861**, that was published in your journal. I intend to use this figure in my Masters Thesis titled: **Six2 EXHIBITS A TEMPORAL-SPATIAL EXPRESSION PROFILE IN THE DEVELOPING MOUSE PALATE AND IMPACTS CELL PROLIFERATION DURING MURINE PALATOGENESIS**. Please let me know if you need more information to process my request. Thank you.

Sincerely,

Dennis.

Dennis Okori Okello
MSc Pharmacy candidate
University of Saskatchewan
S7N 5E5, Saskatoon, SK, CANADA.

RE: Request for permission to use figure

 DELETE  REPLY  REPLY ALL  FORWARD ...

permissions <permissions@biologists.com>

Thu 7/16/2015 4:55 AM

Mark as unread

To: Okello, Dennis Okori; permissions <permissions@biologists.com>;

Dear Dennis,

Thank you for your request. Permission is granted with no charge.

The acknowledgement should state "reproduced / adapted with permission" and give the source journal name - the acknowledgement should either provide full citation details or refer to the relevant citation in the article reference list - the full citation details should include authors, journal, year, volume, issue and page citation.

Where appearing online or in other electronic media, a link should be provided to the original article (e.g. via DOI).

Development: dev.biologists.org

We wish you the best of luck with your thesis.

Kind regards,

Samantha Arnold

Permissions | The Company of Biologists

Bidder Building, Station Road, Histon, Cambridge, CB24 9LF, UK

T: +44 (0) 1223 632867 | samantha.arnold@biologists.com | www.biologists.com

Registered office: The Company Of Biologists Ltd, Bidder Building, Station Road, Histon, Cambridge CB24 9LF, United Kingdom, Registered in England and Wales. Company Limited by Guarantee No 514735. Registered Charity No 277992 The information contained in this message and any attachment is confidential, legally privileged and is intended for the addressee only. Any dissemination, distribution, copying, disclosure or use of this message/attachment or its contents is strictly prohibited and may be unlawful. No contract is intended or implied, unless confirmed by hard copy. If you have received this message in error, please inform the sender and delete it from your mailbox or any other storage mechanism. The Company of Biologists Ltd cannot accept liability for any statements made which are clearly the senders' own and not expressly made on behalf of The



Note: Copyright.com supplies permissions but not the copyrighted content itself.

1
PAYMENT

2
REVIEW

3
CONFIRMATION

Step 3: Order Confirmation

Thank you for your order! A confirmation for your order will be sent to your account email address. If you have questions about your order, you can call us at +1.855.239.3415 Toll Free, M-F between 3:00 AM and 6:00 PM (Eastern), or write to us at info@copyright.com. This is not an invoice.

Confirmation Number: 11400274
Order Date: 07/13/2015

If you paid by credit card, your order will be finalized and your card will be charged within 24 hours. If you choose to be invoiced, you can change or cancel your order until the invoice is generated.

Payment Information

Dennis Okello
deo644@mail.usask.ca
+1 (306) 716-2565
Payment Method: n/a

Order Details

The journal of clinical investigation

Order detail ID: 67703583
Order License Id: 3667310027869
ISSN: 0021-9738
Publication Type: Journal
Volume:
Issue:
Start page:
Publisher: AMERICAN SOCIETY FOR CLINICAL INVESTIGATION
Author/Editor: AMERICAN SOCIETY FOR CLINICAL INVESTIGATION

Permission Status: **Granted**

Permission type: Republish or display content
Type of use: Republish in a thesis/dissertation

Requestor type Academic institution

Format Print, Electronic

Portion chart/graph/table/figure

Number of charts/graphs/tables/figures 1

Title or numeric reference of the portion (s) Figure 1B: Schematic of sections of normal palate shelf development at the indicated days after conception.

Title of the article or chapter the portion is from Murray J.C and Schutte B.C (2004). Cleft palate: players, pathways, and pursuits. J Clin Invest.; 113(12):1676-1678).

Editor of portion(s) N/A

<https://www.copyright.com/printCoiConfirmPurchase.do?operation=defaultOperation&c...> 13/07/2015

Author of portion(s)	N/A
Volume of serial or monograph	113
Issue, if republishing an article from a serial	12
Page range of portion	1676-1678
Publication date of portion	2004
Rights for	Main product
Duration of use	Current edition and up to 5 years
Creation of copies for the disabled	no
With minor editing privileges	yes
For distribution to	Canada
In the following language(s)	Original language of publication
With incidental promotional use	no
Lifetime unit quantity of new product	Up to 499
Made available in the following markets	Educational
The requesting person/organization	Dennis Okello, University of Saskatchewan, S7N 5E5, Saskatoon, SK, CANADA.
Order reference number	
Author/Editor	Jeffrey C. Murray and Brian C. Schutte
The standard identifier	Figure 1B: Schematic of sections of normal palate development at the indicated days after conception
Title	Cleft palate: players, pathways, and pursuits

<https://www.copyright.com/printCoiConfirmPurchase.do?operation=defaultOperation&c...> 13/07/2015

Publisher The Journal of Clinical Investigation

Expected publication date Aug 2015

Estimated size (pages) 140

Note: This item will be invoiced or charged separately through CCC's **RightsLink** service. More info

\$ 0.00

Total order items: 1

This is not an invoice.

Order Total: 0.00 USD

<https://www.copyright.com/printCoiConfirmPurchase.do?operation=defaultOperation&c...> 13/07/2015